

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number  
**WO 03/080027 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 9/14,**  
31/565 (US). **WALTERS, Amy** [US/US]; 5 Shannon Rd., North  
Wales, PA (US).
- (21) International Application Number: **PCT/US03/08546** (74) Agents: **SIMKIN, Michele, M.** et al.; **Foley & Lardner,**  
Washington Harbour, 3000 K Street, N.W., Suite 500,  
Washington, DC 20007-5143 (US).
- (22) International Filing Date: 20 March 2003 (20.03.2003)
- (25) Filing Language: English (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.
- (26) Publication Language: English
- (30) Priority Data:  
60/365,540 20 March 2002 (20.03.2002) US  
60/366,542 25 March 2002 (25.03.2002) US
- (71) Applicant (*for all designated States except US*): **ELAN**  
**PHARMA INTERNATIONAL LTD.** [US/US]; 3000  
Horizon Drive, King of Prussia, PA 19406 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **MERISKO-LIV-**  
**ERSIDGE, Elaine** [US/US]; 258 Colwyn Terrace, West  
Chester, PA 19380 (US). **BOSCH, H., William** [US/US];  
237 Rodney Circle, Bryn Mawr, PA 19010 (US). **CARY,**  
**Greta, G.** [US/US]; 204 Marlbrook Lane, Lansdale, PA  
19446 (US). **PRUITT, John** [US/US]; 603 Buyers Road,  
Collegeville, PA 19426 (US). **RYDE, Tuula** [US/US]; 54  
Lloyd Avenue, Malvern, PA 19355 (US). **JAIN, Rajeev**  
[IN/US]; 121 Honeylocust Ct., Coosooeville, PA 19426
- (84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 03/080027 A1

(54) Title: **NANOPARTICULATE COMPOSITIONS OF ANGIOGENESIS INHIBITORS**

(57) Abstract: Nanoparticulate compositions comprising at least one poorly soluble angiogenesis inhibitor and at least one surface stabilizer are described. The nanoparticulate compositions have an average particle size of less than about 2000 nm. The invention also describes methods of making and using such compositions.

## NANOPARTICULATE COMPOSITIONS OF ANGIOGENESIS INHIBITORS

### FIELD OF THE INVENTION

5           The present invention is directed to nanoparticulate formulations of angiogenesis inhibitors and methods of making and using such compositions.

### BACKGROUND OF THE INVENTION

#### 10    A.     Background Regarding Nanoparticulate Compositions

Nanoparticulate compositions, first described in U.S. Patent No. 5,145,684 ("the '684 patent"), are particles consisting of a poorly soluble therapeutic or diagnostic agent having adsorbed onto the surface thereof a non-crosslinked surface stabilizer. This invention is an improvement over that disclosed in the '684 patent, as the '684 patent  
15   does not describe nanoparticulate compositions comprising an angiogenesis inhibitor.

The '684 patent describes a method of screening active agents to identify useful surface stabilizers that enable the production of a nanoparticulate composition. Not all surface stabilizers will function to produce a stable, non-agglomerated nanoparticulate composition for all active agents.

20       Methods of making nanoparticulate compositions are described in, for example, U.S. Patent Nos. 5,518,187 and 5,862,999, both for "Method of Grinding Pharmaceutical Substances;" U.S. Patent No. 5,718,388, for "Continuous Method of Grinding Pharmaceutical Substances;" and U.S. Patent No. 5,510,118 for "Process of Preparing Therapeutic Compositions Containing Nanoparticles."

25       Nanoparticulate compositions are also described in, for example, U.S. Patent Nos. 5,298,262 for "Use of Ionic Cloud Point Modifiers to Prevent Particle Aggregation During Sterilization;" 5,302,401 for "Method to Reduce Particle Size Growth During Lyophilization;" 5,318,767 for "X-Ray Contrast Compositions Useful in Medical Imaging;" 5,326,552 for "Novel Formulation For Nanoparticulate X-Ray Blood Pool  
30   Contrast Agents Using High Molecular Weight Non-ionic Surfactants;" 5,328,404 for "Method of X-Ray Imaging Using Iodinated Aromatic Propanedioates;" 5,336,507 for

“Use of Charged Phospholipids to Reduce Nanoparticle Aggregation;” 5,340,564 for  
“Formulations Comprising Olin 10-G to Prevent Particle Aggregation and Increase  
Stability;” 5,346,702 for “Use of Non-Ionic Cloud Point Modifiers to Minimize  
Nanoparticulate Aggregation During Sterilization;” 5,349,957 for “Preparation and  
5 Magnetic Properties of Very Small Magnetic-Dextran Particles;” 5,352,459 for “Use of  
Purified Surface Modifiers to Prevent Particle Aggregation During Sterilization;”  
5,399,363 and 5,494,683, both for “Surface Modified Anticancer Nanoparticles;”  
5,401,492 for “Water Insoluble Non-Magnetic Manganese Particles as Magnetic  
Resonance Enhancement Agents;” 5,429,824 for “Use of Tyloxapol as a Nanoparticulate  
10 Stabilizer;” 5,447,710 for “Method for Making Nanoparticulate X-Ray Blood Pool  
Contrast Agents Using High Molecular Weight Non-ionic Surfactants;” 5,451,393 for “X-  
Ray Contrast Compositions Useful in Medical Imaging;” 5,466,440 for “Formulations of  
Oral Gastrointestinal Diagnostic X-Ray Contrast Agents in Combination with  
Pharmaceutically Acceptable Clays;” 5,470,583 for “Method of Preparing Nanoparticle  
15 Compositions Containing Charged Phospholipids to Reduce Aggregation;” 5,472,683 for  
“Nanoparticulate Diagnostic Mixed Carbamic Anhydrides as X-Ray Contrast Agents for  
Blood Pool and Lymphatic System Imaging;” 5,500,204 for “Nanoparticulate Diagnostic  
Dimers as X-Ray Contrast Agents for Blood Pool and Lymphatic System Imaging;”  
5,518,738 for “Nanoparticulate NSAID Formulations;” 5,521,218 for “Nanoparticulate  
20 Iododipamide Derivatives for Use as X-Ray Contrast Agents;” 5,525,328 for  
“Nanoparticulate Diagnostic Diatrizoxy Ester X-Ray Contrast Agents for Blood Pool and  
Lymphatic System Imaging;” 5,543,133 for “Process of Preparing X-Ray Contrast  
Compositions Containing Nanoparticles;” 5,552,160 for “Surface Modified NSAID  
Nanoparticles;” 5,560,931 for “Formulations of Compounds as Nanoparticulate  
25 Dispersions in Digestible Oils or Fatty Acids;” 5,565,188 for “Polyalkylene Block  
Copolymers as Surface Modifiers for Nanoparticles;” 5,569,448 for “Sulfated Non-ionic  
Block Copolymer Surfactant as Stabilizer Coatings for Nanoparticle Compositions;”  
5,571,536 for “Formulations of Compounds as Nanoparticulate Dispersions in Digestible  
Oils or Fatty Acids;” 5,573,749 for “Nanoparticulate Diagnostic Mixed Carboxylic  
30 Anhydrides as X-Ray Contrast Agents for Blood Pool and Lymphatic System Imaging;”

5,573,750 for "Diagnostic Imaging X-Ray Contrast Agents;" 5,573,783 for "Redispersible Nanoparticulate Film Matrices With Protective Overcoats;" 5,580,579 for "Site-specific Adhesion Within the GI Tract Using Nanoparticles Stabilized by High Molecular Weight, Linear Poly(ethylene Oxide) Polymers;" 5,585,108 for "Formulations of Oral

5 Gastrointestinal Therapeutic Agents in Combination with Pharmaceutically Acceptable Clays;" 5,587,143 for "Butylene Oxide-Ethylene Oxide Block Copolymers Surfactants as Stabilizer Coatings for Nanoparticulate Compositions;" 5,591,456 for "Milled Naproxen with Hydroxypropyl Cellulose as Dispersion Stabilizer;" 5,593,657 for "Novel Barium Salt Formulations Stabilized by Non-ionic and Anionic Stabilizers;" 5,622,938 for "Sugar

10 Based Surfactant for Nanocrystals;" 5,628,981 for "Improved Formulations of Oral Gastrointestinal Diagnostic X-Ray Contrast Agents and Oral Gastrointestinal Therapeutic Agents;" 5,643,552 for "Nanoparticulate Diagnostic Mixed Carbonic Anhydrides as X-Ray Contrast Agents for Blood Pool and Lymphatic System Imaging;" 5,718,388 for "Continuous Method of Grinding Pharmaceutical Substances;" 5,718,919 for

15 "Nanoparticles Containing the R(-)Enantiomer of Ibuprofen;" 5,747,001 for "Aerosols Containing Beclomethasone Nanoparticle Dispersions;" 5,834,025 for "Reduction of Intravenously Administered Nanoparticulate Formulation Induced Adverse Physiological Reactions;" 6,045,829 "Nanocrystalline Formulations of Human Immunodeficiency Virus (HIV) Protease Inhibitors Using Cellulosic Surface Stabilizers;" 6,068,858 for "Methods

20 of Making Nanocrystalline Formulations of Human Immunodeficiency Virus (HIV) Protease Inhibitors Using Cellulosic Surface Stabilizers;" 6,153,225 for "Injectable Formulations of Nanoparticulate Naproxen;" 6,165,506 for "New Solid Dose Form of Nanoparticulate Naproxen;" 6,221,400 for "Methods of Treating Mammals Using Nanocrystalline Formulations of Human Immunodeficiency Virus (HIV) Protease

25 Inhibitors;" 6,264,922 for "Nebulized Aerosols Containing Nanoparticle Dispersions;" 6,267,989 for "Methods for Preventing Crystal Growth and Particle Aggregation in Nanoparticle Compositions;" 6,270,806 for "Use of PEG-Derivatized Lipids as Surface Stabilizers for Nanoparticulate Compositions;" 6,316,029 for "Rapidly Disintegrating Solid Oral Dosage Form;" 6,375,986 for "Solid Dose Nanoparticulate Compositions

30 Comprising a Synergistic Combination of a Polymeric Surface Stabilizer and Dioctyl

Sodium Sulfosuccinate," 6,428,814 for "Bioadhesive nanoparticulate compositions having cationic surface stabilizers;" 6,431,478 for "Small Scale Mill;" and 6,432,381 for "Methods for targeting drug delivery to the upper and/or lower gastrointestinal tract," all of which are specifically incorporated by reference. In addition, U.S. Patent Application  
5 No. 20020012675 A1, published on January 31, 2002, for "Controlled Release Nanoparticulate Compositions," describes nanoparticulate compositions, and is specifically incorporated by reference.

Amorphous small particle compositions are described in, for example, U.S. Patent Nos. 4,783,484 for "Particulate Composition and Use Thereof as Antimicrobial Agent;"  
10 4,826,689 for "Method for Making Uniformly Sized Particles from Water-Insoluble Organic Compounds;" 4,997,454 for "Method for Making Uniformly-Sized Particles From Insoluble Compounds;" 5,741,522 for "Ultrasmall, Non-aggregated Porous Particles of Uniform Size for Entrapping Gas Bubbles Within and Methods;" and 5,776,496, for "Ultrasmall Porous Particles for Enhancing Ultrasound Back Scatter."


## 15 **B. Background Regarding Angiogenesis Inhibitors**

Angiogenesis means the formation of new blood vessels. Tumor angiogenesis is the growth of blood vessels from surrounding tissue to a solid tumor, caused by the release  
20 of chemicals by the tumor. Other chemicals, called angiogenesis inhibitors, signal the process to stop. Angiogenesis plays an important role in the growth and spread of cancer, as new blood vessels "feed" the cancer cells with oxygen and nutrients, allowing these cells to grow, invade nearby tissue, spread to other parts of the body, and form new colonies of cancer cells. Because cancer cannot grow or spread without the formation of  
25 new blood vessels, angiogenesis inhibitors can be useful in preventing the growth of cancer by blocking the formation of new blood vessels from surrounding tissue to a solid tumor. This in turn might stop the tumor from growing and spreading to other parts of the body. In animal studies, angiogenesis inhibitors have successfully stopped the formation of new blood vessels, causing the cancer to shrink and die. *See*

30 [http://cis.nci.nih.gov/fact/7\\_42.htm](http://cis.nci.nih.gov/fact/7_42.htm).

Exemplary angiogenesis inhibitors given at cancer.gov (affiliated with the National Institutes of Health) are provided in the following table.

Agent	Description
2-methoxyestradiol	2ME. A drug derived from estrogen that belongs to the family of drugs called angiogenesis inhibitors. It prevents the formation of new blood vessels that tumors need to grow.
AG3340	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors. AG3340 is a matrix metalloproteinase (MMP) inhibitor. Also called prinomastat.
batimastat	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors. Batimastat is a matrix metalloproteinase inhibitor.
BAY 12-9566	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
carboxyamidotriazole	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
CC-1088	A drug that is similar but not identical to thalidomide and is being studied as an anticancer drug. It belongs to the family of drugs called angiogenesis inhibitors.
dextromethorphan acetic acid	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
dimethylxanthenone acetic acid	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
EMD 121974	A substance that is being studied as an anticancer and antiangiogenesis drug.
endostatin	A drug that is being studied for its ability to prevent the growth of new blood vessels into a solid tumor. Endostatin belongs to the family of drugs called angiogenesis inhibitors.
IM-862	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
marimastat	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors. Marimastat is a MMP inhibitor.
matrix metalloproteinase	A member of a group of enzymes that can break down proteins, such as collagen, that are normally found in the spaces between cells in tissues (i.e., extracellular matrix proteins). Because these enzymes need zinc or calcium atoms to work properly, they are called metalloproteinases. Matrix metalloproteinases are involved in wound healing, angiogenesis, and tumor cell metastasis.
penicillamine	A drug that removes copper from the body and is used to treat diseases in which there is an excess of this metal. It is also being studied as a possible angiogenesis inhibitor in brain tumors.
PTK787/ZK 222584	An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors.
RPI.4610	A substance that is being studied as a treatment for cancer. It belongs to the family of drugs called angiogenesis inhibitors.
squalamine lactate	A drug that belongs to the family of drugs called angiogenesis inhibitors. It prevents the growth of new blood vessels into a solid tumor.

Agent	Description
SU5416	<p>An anticancer drug that belongs to the family of drugs called angiogenesis inhibitors. SU5416, 3-[2, 4-dimethylpyrrol-5-yl methylidenyl]-2-indolinone, has the following structure <a href="http://www.pharmquest.com/source/features/AAPS_Trends_eRD/SUGEN_Arun_Koparkar.pdf">http://www.pharmquest.com/source/features/AAPS_Trends_eRD/SUGEN_Arun_Koparkar.pdf</a>:</p> 
thalidomide	A drug that belongs to the family of drugs called angiogenesis inhibitors. It prevents the growth of new blood vessels into a solid tumor.
TNP-470	A drug that belongs to the family of drugs called angiogenesis inhibitors. It prevents the growth of new blood vessels into a solid tumor.

Other known angiogenesis inhibitors include, but are not limited to, suramin, combretastatin, paclitaxel, and tamoxifen,. One of these compounds, suramin, is soluble  
5 in water. More detailed descriptions of select angiogenesis inhibitors are given below.

Combretastatin was disclosed in the *Journal of the National Cancer Institute* on April 5, 2000, as an angiogenesis inhibitor isolated from the bark of a South African species of willow tree. The compound is described and claimed in U.S. Patent No. 4,996,237, assigned to the Arizona Board of Regents.

10 2-methoxyestradiol was disclosed in the *Journal of the National Cancer Institute* on April 5, 2000, as an angiogenesis inhibitor. In a press release of February 14, 2000, Entremed, Inc., in Rockville, MD was given permission for Phase I trials of 2ME2. Entremed provides an overview of 2ME2 on their web site. Claim 2 of U.S. Patent No. 5,504,074 is directed to a method for treating mammalian disease characterized by  
15 undesirable angiogenesis comprising administering 2-methoxyestradiol.

At the 54<sup>th</sup> meeting of the Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research, Division of Oncology, the director of the Angiogenesis Foundation informed the committee about the angiogenesis inhibitory activity of paclitaxel. The Merck Index listing of Taxol

(trademark name of paclitaxel) states that the compound was first isolated from the bark of the Pacific yew tree.

At the 58<sup>th</sup> meeting of the Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research, Oncologic Drugs Advisory Committee, it was reported that tamoxifen is an angiogenesis inhibitor. Conventional tamoxifen is generic, as its isolation and identification were described in the 1960s. However, isomers of tamoxifen are patented. *See e.g.*, claim 2 of U.S. Patent No. 4,536,516,.

Newton, "Novel Chemotherapeutic Agents for the Treatment of Brain Cancer," *Expert Opin. Investigational Drugs*, 9:2815-29 (2000), discloses that neoplastic angiogenesis and brain tumor invasion are also targets for therapeutic interventions with new agents such as thalidomide, suramin, and marimastat.

Liekens et al., "Angiogenesis: Regulators and Clinical Applications," *Biochem. Pharmacol.*, 61i:253-70 (2001), disclose that TNP-470 is an angiogenesis inhibitor.

Claim 1 of U.S. Patent No. 5,166,172, assigned to Takeda Chemical Industries, Ltd., is directed to O-(chloroacetylcarbonyl) fumagillol (TNP-470). Example 8 of this patent discloses that TNP-470 is obtained from silica gel with a mixture of n-hexane and ethylacetate.

Experiments examining thalidomide's enantiomers reveal that the S(-) enantiomer has the strongest antiangiogenic activity. Kenyon et al., "Effects of thalidomide and related metabolites in a mouse corneal model of neovascularization," *Exp. Eye Res.*, 64:971-978 (1997). Moreover, the immunomodulating and anti-inflammatory effects of thalidomide are likely chiefly exerted by S-thalidomide. Eriksson et al., "Intravenous formulations of the enantiomers of thalidomide: Pharmacokinetic and initial pharmacodynamic characterization in man," *J. Pharm. Pharmacol.*, 52:807-817 (2000).

Other studies have shown that the R-isomer provides the drug's sedative effect, and that the S-isomer is responsible for the birth defects associated with the agent. C. Star, "Splitting pairs: molecular maneuver aims for better drugs," *Drug Topics*, 136(15):26 (Aug. 3, 1992).



U.S. Patent No. 6,124,322 teaches that pure enantiomers of thalidomide are converted back into the racemate *in vitro* and *in vivo*. See also *Drug Topics*, above. The antipode is formed immediately after the parenteral administration of one of the isomers of thalidomide *in vivo*, and an equilibrium is established after about 4 hours.

5       The claims of U.S. Patent No. 6,124,322 recite aqueous thalidomide solutions of either the R or S enantiomers of thalidomide. According to the disclosure of the patent, the enantiomers are more soluble than the racemate of thalidomide, which enables intravenous administration of the enantiomers.

Angiogenesis inhibitors currently in clinical trials include the following  
10   ([http://www.cancer.gov/clinical\\_trials/doc.aspx?viewid=B0959CBB-3004-4160-A679-6DD204BEE68C](http://www.cancer.gov/clinical_trials/doc.aspx?viewid=B0959CBB-3004-4160-A679-6DD204BEE68C)): marimastat, COL-3 (synthetic MMP inhibitor; tetracycline derivative), neovastat (naturally occurring MMP inhibitor), BMS-275291 (synthetic MMP inhibitor), thalidomide, squalamine (extract from dogfish shark liver; inhibits sodium-hydrogen exchanger, NHE3), 2-ME (inhibition of endothelial cells), SU6668 (blocks  
15   VEGF, FGF, and PDGF receptor signaling), interferon-alpha (inhibition of bFGF and VEGF production), anti-VEGF antibody (monoclonal antibody to vascular endothelial growth factor (VEGF)), Medi-522 (Vitaxin II) (antibody that blocks the integrin present on endothelial cell surface), EMD121974 (small molecule blocker of integrin present on endothelial cell surface), CAI (inhibitor of calcium influx), celecoxib (enzyme cyclo-  
20   oxygenase 2 (COX-2)), Interleukin-12 (up-regulation of interferon gamma and IP-10), and IM862 (unknown mechanism).

Additionally, the following angiogenesis inhibitors are disclosed in the CalBioChem® catalog at page xxxiii: Amiloride, Human Angiostatin® Protein, Human Angiostatin K1-3, Human Angiostatin K1-5, Captopril, DL-alpha-  
25   Difluoromethylornithine HCl, Human Recombinant Endostatin™ Protein (*Pichia pastoris*), Mouse Recombinant Endostatin™ Protein (*Pichia pastoris*), Mouse Recombinant His-Tag® Endostatin™ Protein (*Spodoptera frugiperda*), Fumagillin (*Aspergillus fumigatus*), Herbimycin A (*Streptomyces* sp), 4-Hydroxyphenylretinamide, Mouse Recombinant alpha-interferon (*E. coli*), Human Recombinant gamma-interferon  
30   (*E. coli*), Juglone, Laminin Hexapeptide, Laminin Pentapeptide, Lavendustin A,

Medroxyprogesterone Acetate, 2-Methoxyestradiol, Minocycline HCl, Human  
Recombinant Placental Ribonuclease Inhibitor, Sodium Salt Suramin, (+)-Thalidomide  
Human Platelet Thrombospondin, Recombinant Bovine Tissue Inhibitor of  
Metalloproteinase 1, Recombinant Human Tissue Inhibitor of Metalloproteinase 1,  
5 Recombinant Human Neutrophil Granulocyte Tissue Inhibitor of Metalloproteinase 1, and  
Recombinant Human Rheumatoid Synovial Fibroblast Tissue Inhibitor of  
Metalloproteinase 2.

\* \* \* \*

10 There is a need in the art for nanoparticulate compositions of angiogenesis  
inhibitors and methods of making and using such compositions. The present invention  
satisfies these needs.

#### **SUMMARY OF THE INVENTION**

15 The present invention is directed to nanoparticulate compositions comprising at  
least one poorly soluble angiogenesis inhibitor and at least one surface stabilizer  
associated with the surface of the angiogenesis inhibitor.

20 Another aspect of the invention is directed to pharmaceutical compositions  
comprising a nanoparticulate angiogenesis inhibitor composition of the invention. The  
pharmaceutical compositions preferably comprise at least one poorly soluble angiogenesis  
inhibitor, at least one surface stabilizer associated with the surface of the inhibitor, and a  
pharmaceutically acceptable carrier, as well as any desired excipients.

25 This invention further discloses a method of making a nanoparticulate  
composition having at least one poorly soluble angiogenesis inhibitor and at least one  
surface stabilizer associated with the surface of the inhibitor. Such a method comprises  
contacting a poorly soluble nanoparticulate angiogenesis inhibitor with at least one  
surface stabilizer for a time and under conditions sufficient to provide an angiogenesis  
inhibitor/surface stabilizer composition. The surface stabilizer can be contacted with the  
30 angiogenesis inhibitor either before, during, or after particle size reduction of the  
angiogenesis inhibitor.

The present invention is further directed to a method of treatment comprising administering to a mammal a therapeutically effective amount of a nanoparticulate angiogenesis inhibitor composition according to the invention.

Both the foregoing general description and the following detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed. Other objects, advantages, and novel features will be readily apparent to those skilled in the art from the following detailed description of the invention.

#### **DETAILED DESCRIPTION OF THE INVENTION**

The present invention is directed to the surprising and unexpected discovery that stable nanoparticulate compositions of angiogenesis inhibitors can be made.

Advantages of the angiogenesis inhibitor compositions of the invention include, but are not limited to: (1) faster onset of action; (2) smaller tablet or other solid dosage form size, or smaller volume if in a liquid dosage form; (3) smaller doses of drug required to obtain the same pharmacological effect as compared to conventional microcrystalline forms of the same angiogenesis inhibitor; (4) increased bioavailability as compared to conventional microcrystalline forms of the same angiogenesis inhibitor; (5) substantially similar pharmacokinetic profiles of the angiogenesis inhibitor compositions of the invention when administered in the fed versus the fasted state; (6) bioequivalency of the angiogenesis inhibitor compositions of the invention when administered in the fed versus the fasted state; (7) improved pharmacokinetic profiles; (8) an increased rate of dissolution for the angiogenesis inhibitor compositions of the invention as compared to conventional microcrystalline forms of the same angiogenesis inhibitor; (9) bioadhesive angiogenesis inhibitor compositions; (10) the angiogenesis inhibitor compositions of the invention can be sterile filtered; and (11) the angiogenesis inhibitor compositions of the invention can be used in conjunction with other active agents.

The invention encompasses the angiogenesis inhibitor compositions of the invention formulated or coadministered with one or more non-angiogenesis inhibitor

active agents, either conventional (solubilized or microparticulate) or nanoparticulate. Methods of using such combination compositions are also encompassed by the invention.

The present invention is described herein using several definitions, as set forth below and throughout the application.

5        "About" will be understood by persons of ordinary skill in the art and will vary to some extent on the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

10        As used herein with reference to stable drug particles, 'stable' means that angiogenesis inhibitor particles do not appreciably flocculate or agglomerate due to interparticle attractive forces or otherwise increase in particle size.

15        "Therapeutically effective amount" as used herein with respect to a drug dosage, shall mean that dosage that provides the specific pharmacological response for which the drug is administered in a significant number of subjects in need of such treatment. It is emphasized that 'therapeutically effective amount,' administered to a particular subject in a particular instance will not always be effective in treating the diseases described herein, even though such dosage is deemed a 'therapeutically effective amount' by those skilled in the art. It is to be further understood that drug dosages are, in particular instances, measured as oral dosages, or with reference to drug levels as measured in blood.

20        "Conventional active agents or drugs" refers to non-nanoparticulate or solubilized active agents or drugs. Non-nanoparticulate active agents have an effective average particle size of greater than about 2 microns.

**A. Preferred Characteristics of the Angiogenesis Inhibitor Compositions of the Invention**

**1. Fast Onset of Activity**

5 The use of conventional formulations of angiogenesis inhibitors is not ideal due to delayed onset of action. In contrast, the nanoparticulate angiogenesis inhibitor compositions of the invention exhibit faster therapeutic effects. Moreover, nanoparticulate formulations of angiogenesis inhibitors enable selection of an  
10 angiogenesis inhibitor with a long half-life in the blood stream while still providing the subject with a fast-acting compound.

Preferably, following administration the angiogenesis inhibitor compositions of the invention have a  $T_{max}$  of less than about 2.5 hours, less than about 2.25 hours, less than about 2 hours, less than about 1.75 hours, less than about 1.5 hours, less than about  
15 1.25 hours, less than about 1.0 hours, less than about 50 minutes, less than about 40 minutes, less than about 30 minutes, less than about 25 minutes, less than about 20 minutes, less than about 15 minutes, or less than about 10 minutes.

**2. Increased Bioavailability**

20 The angiogenesis inhibitor compositions of the invention preferably exhibit increased bioavailability, at the same dose of the same angiogenesis inhibitor, and require smaller doses, as compared to prior conventional angiogenesis inhibitor compositions.

Any drug, including angiogenesis inhibitors, can have adverse side effects. Thus, lower doses of angiogenesis inhibitors which can achieve the same or better therapeutic  
25 effects as those observed with larger doses of conventional angiogenesis inhibitors are desired. Such lower doses can be realized with the angiogenesis inhibitor compositions of the invention, because the greater bioavailability observed with the nanoparticulate angiogenesis inhibitor compositions as compared to conventional drug formulations means that smaller doses of drug are required to obtain the desired therapeutic effect.

**3. The Pharmacokinetic Profiles of the Angiogenesis Inhibitor Compositions of the Invention are not Substantially Affected by the Fed or Fasted State of the Subject Ingesting the Compositions**

5 The invention encompasses an angiogenesis inhibitor composition wherein the pharmacokinetic profile of the angiogenesis inhibitor is not substantially affected by the fed or fasted state of a subject ingesting the composition. This means that there is no substantial difference in the quantity of drug absorbed or the rate of drug absorption when  
10 the nanoparticulate angiogenesis inhibitor compositions are administered in the fed versus the fasted state. Thus, the nanoparticulate angiogenesis inhibitor compositions of the invention substantially eliminate the effect of food on the pharmacokinetics of the angiogenesis inhibitor.

15 Preferably, the difference in absorption of the nanoparticulate angiogenesis inhibitor compositions of the invention, when administered in the fed versus the fasted state, is less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 35%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, less than about 3%, or essentially no  
20 difference.

In addition, preferably the difference in the rate of absorption (*i.e.*,  $T_{max}$ ) of the nanoparticulate angiogenesis inhibitor compositions of the invention, when administered in the fed versus the fasted state, is less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than  
25 about 40%, less than about 30%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, less than about 3%, or essentially no difference.

Benefits of a dosage form which substantially eliminates the effect of food include an increase in subject convenience, thereby increasing subject compliance, as the subject does not need to ensure that they are taking a dose either with or without food.

#### 4. Redispersibility Profiles of the Angiogenesis Inhibitor Compositions of the Invention

An additional feature of the angiogenesis inhibitor compositions of the invention is that the compositions redisperse such that the effective average particle size of the redispersed angiogenesis inhibitor particles is less than about 2 microns. This is significant, as if upon administration the nanoparticulate angiogenesis inhibitor compositions of the invention did not redisperse to a substantially nanoparticulate particle size, then the dosage form may lose the benefits afforded by formulating the angiogenesis inhibitor into a nanoparticulate particle size.

This is because nanoparticulate angiogenesis inhibitor compositions benefit from the small particle size of the angiogenesis inhibitor; if the nanoparticulate angiogenesis inhibitor particles do not redisperse into the small particle sizes upon administration, then "clumps" or agglomerated angiogenesis inhibitor particles are formed, owing to the extremely high surface free energy of the nanoparticulate system and the thermodynamic driving force to achieve an overall reduction in free energy. With the formation of such agglomerated particles, the bioavailability of the dosage form may fall well below that observed with the liquid dispersion form of the nanoparticulate angiogenesis inhibitor composition.

Preferably, the redispersed angiogenesis inhibitor particles of the invention have an effective average particle size of less than about 2 microns, less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 75 nm, or less than about 50 nm, as measured by light-scattering methods, microscopy, or other appropriate methods.

## 5. Bioadhesive Angiogenesis Inhibitor Compositions

Bioadhesive angiogenesis inhibitor compositions of the invention comprise at least one cationic surface stabilizer, which are described in more detail below.

Bioadhesive formulations of angiogenesis inhibitors exhibit exceptional bioadhesion to

5 biological surfaces, such as mucous. The term bioadhesion refers to any attractive interaction between two biological surfaces or between a biological and a synthetic surface. In the case of bioadhesive nanoparticulate angiogenesis inhibitor compositions, the term bioadhesion is used to describe the adhesion between the nanoparticulate angiogenesis inhibitor compositions and a biological substrate (*i.e.* gastrointestinal mucin,  
10 lung tissue, nasal mucosa, *etc.*). *See e.g.*, U.S. Patent No. 6,428,814 for "Bioadhesive Nanoparticulate Compositions Having Cationic Surface Stabilizers," which is specifically incorporated by reference.

The bioadhesive angiogenesis inhibitor compositions of the invention are useful in any situation in which it is desirable to apply the compositions to a biological surface.

15 The bioadhesive angiogenesis inhibitor compositions coat the targeted surface in a continuous and uniform film which is invisible to the naked human eye.

A bioadhesive angiogenesis inhibitor composition slows the transit of the composition, and some angiogenesis inhibitor particles would also most likely adhere to tissue other than the mucous cells and therefore give a prolonged exposure to the  
20 angiogenesis inhibitor, thereby increasing absorption and the bioavailability of the administered dosage.

## 6. Pharmacokinetic Profiles of the Angiogenesis Inhibitor Compositions of the Invention

25

The present invention provides compositions of one or more angiogenesis inhibitors having a desirable pharmacokinetic profile when administered to mammalian subjects. Preferably, the  $T_{max}$  of an administered dose of a nanoparticulate angiogenesis inhibitor is less than that of a conventional non-nanoparticulate composition of the same  
30 angiogenesis inhibitor, administered at the same dosage. In addition, preferably the  $C_{max}$  of a nanoparticulate composition of an angiogenesis inhibitor is greater than the  $C_{max}$  of a



conventional non-nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage.

In comparative pharmacokinetic testing with a non-nanoparticulate composition of an angiogenesis inhibitor, a nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage, preferably exhibits a  $T_{max}$  which is less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, or less than about 10% of the  $T_{max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

In comparative pharmacokinetic testing with a non-nanoparticulate composition of an angiogenesis inhibitor, a nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage, preferably exhibits a  $C_{max}$  which is greater than about 5%, greater than about 10%, greater than about 15%, greater than about 20%, greater than about 30%, greater than about 40%, greater than about 50%, greater than about 60%, greater than about 70%, greater than about 80%, greater than about 90%, greater than about 100%, greater than about 110%, greater than about 120%, greater than about 130%, greater than about 140%, or greater than about 150% than the  $C_{max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

The desirable pharmacokinetic profile, as used herein, is the pharmacokinetic profile measured after an initial dose of an angiogenesis inhibitor. The compositions can be formulated in any way as described below.

### C. Combination Pharmacokinetic Profile Compositions

In yet another embodiment of the invention, a first angiogenesis inhibitor composition providing a desired pharmacokinetic profile is co-administered, sequentially administered, or combined with at least one other angiogenesis inhibitor composition that generates a desired different pharmacokinetic profile. More than two angiogenesis inhibitor compositions can be co-administered, sequentially administered, or combined. While at least one of the angiogenesis inhibitor compositions has a nanoparticulate

particle size, the additional one or more angiogenesis inhibitor compositions can be nanoparticulate, solubilized, or have a conventional microparticulate particle size.

For example, a first angiogenesis inhibitor composition can have a nanoparticulate particle size, conferring a short  $T_{max}$  and typically a higher  $C_{max}$ . This first angiogenesis inhibitor composition can be combined, co-administered, or sequentially administered with a second composition comprising: (1) a different nanoparticulate angiogenesis inhibitor exhibiting slower absorption and, therefore a longer  $T_{max}$  and typically a lower  $C_{max}$ ; (2) the same angiogenesis inhibitor having a larger (but still nanoparticulate) particle size, and therefore exhibiting slower absorption, a longer  $T_{max}$ , and typically a lower  $C_{max}$ ; or (3) a microparticulate angiogenesis inhibitor composition (with the angiogenesis inhibitor being either the same as or different from the angiogenesis inhibitor of the first composition), exhibiting a longer  $T_{max}$ , and typically a lower  $C_{max}$ .

The second, third, fourth, *etc.*, angiogenesis inhibitor composition can differ from the first, and from each other, for example: (1) in the identity of the angiogenesis inhibitor; (2) in the effective average particle sizes of each composition; or (3) in the dosage of the angiogenesis inhibitor. Angiogenesis inhibitor compositions can produce a different  $T_{max}$ . Such a combination composition can reduce the dose frequency required.

If the second angiogenesis inhibitor composition has a nanoparticulate particle size, then preferably the angiogenesis inhibitor has at least one surface stabilizer associated with the surface of the drug particles. The one or more surface stabilizers can be the same as or different from the surface stabilizers associated with the surface of the first angiogenesis inhibitor.

Preferably where co-administration of a "fast-acting" formulation and a "longer-lasting" formulation is desired, the two formulations are combined within a single composition, for example a dual-release composition.

#### D. Compositions

The compositions of the invention comprise at least one poorly soluble angiogenesis inhibitor and at least one surface stabilizer. Surface stabilizers useful herein associate with the surface of the nanoparticulate angiogenesis inhibitor, but do not

chemically react with the angiogenesis inhibitor or itself. Preferably, individually adsorbed molecules of the surface stabilizer are essentially free of intermolecular cross-linkages.

The present invention also includes nanoparticulate angiogenesis inhibitors having at least one surface stabilizer associated with the surface thereof, formulated into compositions together with one or more non-toxic physiologically acceptable carriers, adjuvants, or vehicles, collectively referred to as carriers.

### 1. Angiogenesis Inhibitor Drug Particles

The compositions of the invention comprise a poorly soluble angiogenesis inhibitor which is dispersible in at least one liquid medium. The angiogenesis inhibitor exists as a discrete crystalline phase, as an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, or a combination thereof. The crystalline phase differs from a non-crystalline or amorphous phase which results from precipitation techniques, such as those described in EP Patent No. 275,796. By "poorly soluble" it is meant that the angiogenesis inhibitor has a solubility in a liquid dispersion medium of less than about 30 mg/mL, less than about 20 mg/mL, less than about 10 mg/mL, or less than about 1 mg/mL. Useful liquid dispersion mediums include, but are not limited to, water, aqueous salt solutions, safflower oil, and solvents such as ethanol, t-butanol, hexane, and glycol.

Useful angiogenesis inhibitors according to the invention include, but are not limited to: 2-methoxyestradiol, prinomastat, batimastat, BAY 12-9566, carboxyamidotriazole, CC-1088, dextromethorphan acetic, dimethylxanthenone acetic acid, EMD 121974, endostatin, IM-862, marimastat, matrix metalloproteinase, penicillamine, PTK787/ZK 222584, RPI.4610, squalamine, squalamine lactate, SU5416, (+)-thalidomide, S- thalidomide, R- thalidomide, TNP-470, combretastatin, paclitaxel, tamoxifen, COL-3, neovastat, BMS-275291, SU6668, interferon-alpha, anti-VEGF antibody, Medi-522 (Vitaxin II), CAI, celecoxib, Interleukin-12, IM862, Amiloride, Angiostatin® Protein, Angiostatin K1-3, Angiostatin K1-5, Captopril, DL-alpha-Difluoromethylornithine, DL-alpha-Difluoromethylornithine HCl, His-Tag® Endostatin™ Protein, Fumagillin, Herbimycin A, 4-Hydroxyphenylretinamide, gamma-interferon,

Juglone, Laminin, Laminin Hexapeptide, Laminin Pentapeptide, Lavendustin A, Medroxyprogesterone, Medroxyprogesterone Acetate, Minocycline, Minocycline HCl, Placental Ribonuclease Inhibitor, Suramin, Sodium Salt Suramin, Human Platelet Thrombospondin, Tissue Inhibitor of Metalloproteinase 1, Neutrophil Granulocyte Tissue Inhibitor of Metalloproteinase 1, and Rheumatoid Synovial Fibroblast Tissue Inhibitor of Metalloproteinase 2. See [http://cis.nci.nih.gov/fact/7\\_42.htm](http://cis.nci.nih.gov/fact/7_42.htm); CalBioChem® catalog at page xxxiii; and [http://www.cancer.gov/clinical\\_trials/doc.aspx?viewid=B0959CBB-3004-4160-A679-6DD204BEE68C](http://www.cancer.gov/clinical_trials/doc.aspx?viewid=B0959CBB-3004-4160-A679-6DD204BEE68C).

## 2. Non-Angiogenesis Inhibitor Active Agents

The nanoparticulate angiogenesis inhibitor compositions of the invention can additionally comprise one or more non-angiogenesis inhibitor active agents, in either a conventional or nanoparticulate particle size. The non-angiogenesis inhibitor active agents can be present in a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, or a mixture thereof.

If the non-angiogenesis inhibitor active agent has a nanoparticulate particle size *i.e.*, a particle size of less than about 2 microns, then preferably it will have one or more surface stabilizers associated with the surface of the active agent. In addition, if the active agent has a nanoparticulate particle size, then it is preferably poorly soluble and dispersible in at least one liquid dispersion medium. By "poorly soluble" it is meant that the active agent has a solubility in a liquid dispersion medium of less than about 30 mg/mL, less than about 20 mg/mL, less than about 10 mg/mL, or less than about 1 mg/mL. Useful liquid dispersion mediums include, but are not limited to, water, aqueous salt solutions, safflower oil, and solvents such as ethanol, t-butanol, hexane, and glycol.

Such active agents can be, for example, a therapeutic agent. A therapeutic agent can be a pharmaceutical agent, including biologics such as amino acids, proteins, peptides, and nucleotides. The active agent can be selected from a variety of known classes of drugs, including, for example, amino acids, proteins, peptides, nucleotides, anti-obesity drugs, central nervous system stimulants, carotenoids, corticosteroids, elastase inhibitors, anti-fungals, oncology therapies, anti-emetics, analgesics, cardiovascular

agents, anti-inflammatory agents, such as NSAIDs and COX-2 inhibitors, anthelmintics, anti-arrhythmic agents, antibiotics (including penicillins), anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives (hypnotics and neuroleptics), astringents, alpha-adrenergic receptor blocking agents, beta-adrenoceptor blocking agents, blood products and substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants (expectorants and mucolytics), diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics (antiparkinsonian agents), haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio-pharmaceuticals, sex hormones (including steroids), anti-allergic agents, stimulants and anoretics, sympathomimetics, thyroid agents, vasodilators, and xanthines.

A description of these classes of active agents and a listing of species within each class can be found in Martindale's *The Extra Pharmacopoeia*, 31<sup>st</sup> Edition (The Pharmaceutical Press, London, 1996), specifically incorporated by reference. The active agents are commercially available and/or can be prepared by techniques known in the art.

Exemplary nutraceuticals and dietary supplements are disclosed, for example, in Roberts et al., *Nutraceuticals: The Complete Encyclopedia of Supplements, Herbs, Vitamins, and Healing Foods* (American Nutraceutical Association, 2001), which is specifically incorporated by reference. Dietary supplements and nutraceuticals are also disclosed in *Physicians' Desk Reference for Nutritional Supplements*, 1st Ed. (2001) and *The Physicians' Desk Reference for Herbal Medicines*, 1st Ed. (2001), both of which are also incorporated by reference. A nutraceutical or dietary supplement, also known as phytochemicals or functional foods, is generally any one of a class of dietary supplements, vitamins, minerals, herbs, or healing foods that have medical or pharmaceutical effects on the body.

Exemplary nutraceuticals or dietary supplements include, but are not limited to, lutein, folic acid, fatty acids (e.g., DHA and ARA), fruit and vegetable extracts, vitamin and mineral supplements, phosphatidylserine, lipoic acid, melatonin,

glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids (*e.g.*, arginine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine), green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish and marine animal oils, and probiotics. Nutraceuticals and dietary supplements also include bio-engineered foods genetically engineered to have a desired property, also known as "pharmafoods."

The compound to be administered in combination with a nanoparticulate angiogenesis inhibitor composition of the invention can be formulated separately from the angiogenesis inhibitor composition or co-formulated with the angiogenesis inhibitor composition. Where an angiogenesis inhibitor composition is co-formulated with a second active agent, the second active agent can be formulated in any suitable manner, such as immediate-release, rapid-onset, sustained-release, or dual-release form.

### 3. Surface Stabilizers

Useful surface stabilizers, which are known in the art and described in the '684 patent, are believed to include those which associate with the surface of the angiogenesis inhibitor but do not chemically bond to or interact with the angiogenesis inhibitor. The surface stabilizer is associated with the surface of the angiogenesis inhibitor in an amount sufficient to maintain the angiogenesis inhibitor particles at an effective average particle size of less than about 2000 nm. Furthermore, the individually adsorbed molecules of the surface stabilizer are preferably essentially free of intermolecular cross-linkages. Two or more surface stabilizers can be employed in the compositions and methods of the invention.

Suitable surface stabilizers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products, and surfactants. Surface stabilizers include nonionic, cationic, zwitterionic, and ionic surfactants.

Representative examples of surface stabilizers include gelatin, casein, lecithin (phosphatides), dextran, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol

emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers (*e.g.*, macrogol ethers such as cetomacrogol 1000), polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters (*e.g.*, the commercially available Tweens<sup>®</sup> such as *e.g.*, Tween 20<sup>®</sup> and Tween 80<sup>®</sup> (ICI Speciality Chemicals)); polyethylene glycols (*e.g.*, Carbowaxs 3550<sup>®</sup> and 934<sup>®</sup> (Union Carbide)), polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose phthalate, noncrystalline cellulose, magnesium aluminium silicate, triethanolamine, polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol, superione, and triton), poloxamers (*e.g.*, Pluronic F68<sup>®</sup> and F108<sup>®</sup>, which are block copolymers of ethylene oxide and propylene oxide); poloxamines (*e.g.*, Tetronic 908<sup>®</sup>, also known as Poloxamine 908<sup>®</sup>, which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF Wyandotte Corporation, Parsippany, N.J.)); Tetronic 1508<sup>®</sup> (T-1508) (BASF Wyandotte Corporation), dialkylesters of sodium sulfosuccinic acid (*e.g.*, Aerosol OT<sup>®</sup>, which is a dioctyl ester of sodium sulfosuccinic acid (DOSS) (American Cyanamid)); Duponol P<sup>®</sup>, which is a sodium lauryl sulfate (DuPont); Tritons X-200<sup>®</sup>, which is an alkyl aryl polyether sulfonate (Rohm and Haas); Crodestas F-110<sup>®</sup>, which is a mixture of sucrose stearate and sucrose distearate (Croda Inc.); p-isononylphenoxypoly-(glycidol), also known as Olin-LOG<sup>®</sup> or Surfactant 10-G<sup>®</sup> (Olin Chemicals, Stamford, CT); Crodestas SL-40<sup>®</sup> (Croda, Inc.); and SA9OHCO, which is C<sub>18</sub>H<sub>37</sub>CH<sub>2</sub>(CON(CH<sub>3</sub>)-CH<sub>2</sub>(CHOH)<sub>4</sub>(CH<sub>2</sub>OH)<sub>2</sub> (Eastman Kodak Co.); decanoyl-N-methylglucamide; n-decyl β-D-glucopyranoside; n-decyl β-D-maltopyranoside; n-dodecyl β-D-glucopyranoside; n-dodecyl β-D-maltoside; heptanoyl-N-methylglucamide; n-heptyl-β-D-glucopyranoside; n-heptyl β-D-thioglucoside; n-hexyl β-D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl β-D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl-β-D-glucopyranoside; octyl β-D-thioglucopyranoside; PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin A, PEG-vitamin E, lysozyme, random copolymers of vinyl pyrrolidone and vinyl acetate, and the like.

Examples of useful cationic surface stabilizers include, but are not limited to, polymers, biopolymers, polysaccharides, cellulose, alginates, phospholipids, and nonpolymeric compounds, such as zwitterionic stabilizers, poly-n-methylpyridinium, anthryl pyridinium chloride, cationic phospholipids, chitosan, polylysine, polyvinylimidazole, polybrene, polymethylmethacrylate trimethylammoniumbromide bromide (PMMTMABr), hexyldesyltrimethylammonium bromide (HDMAB), and polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate dimethyl sulfate.

Other useful cationic stabilizers include, but are not limited to, cationic lipids, sulfonium, phosphonium, and quaternary ammonium compounds, such as stearyltrimethylammonium chloride, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride or bromide, coconut methyl dihydroxyethyl ammonium chloride or bromide, decyl triethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride or bromide, C<sub>12-15</sub>dimethyl hydroxyethyl ammonium chloride or bromide, coconut dimethyl hydroxyethyl ammonium chloride or bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride or bromide, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride or bromide, N-alkyl (C<sub>12-18</sub>)dimethylbenzyl ammonium chloride, N-alkyl (C<sub>14-18</sub>)dimethyl-benzyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and (C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride, trimethylammonium halide, alkyl-trimethylammonium salts and dialkyl-dimethylammonium salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt and/or an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyldimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride and dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C<sub>12</sub>, C<sub>15</sub>, C<sub>17</sub> trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl ammonium chlorides, alkyldimethylammonium halogenides, tricetyl methyl ammonium chloride,



decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride (ALQUAT 336™), POLYQUAT 10™ (polyquaternium 10; Buckman Laboratories, TN), tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters (such as choline esters of fatty acids), benzalkonium chloride, stearylalkonium chloride compounds (such as stearyltrimonium chloride and Di-stearyldimonium chloride), cetyl pyridinium bromide or chloride, halide salts of quaternized polyoxyethylalkylamines, MIRAPOL™ (quaternized ammonium salt polymers) and ALKAQUAT™ (benzalkonium chloride) (Alkaril Chemical Company), alkyl pyridinium salts; amines, such as alkylamines, dialkylamines, alkanolamines, polyethylenepolyamines, N,N-dialkylaminoalkyl acrylates, and vinyl pyridine, amine salts, such as lauryl amine acetate, stearyl amine acetate, alkylpyridinium salt, and alkylimidazolium salt, and amine oxides; imide azolinium salts; protonated quaternary acrylamides; methylated quaternary polymers, such as poly[diallyl dimethylammonium chloride] and poly-[N-methyl vinyl pyridinium chloride]; and cationic guar.

Such exemplary cationic surface stabilizers and other useful cationic surface stabilizers are described in J. Cross and E. Singer, *Cationic Surfactants: Analytical and Biological Evaluation* (Marcel Dekker, 1994); P. and D. Rubingh (Editor), *Cationic Surfactants: Physical Chemistry* (Marcel Dekker, 1991); and J. Richmond, *Cationic Surfactants: Organic Chemistry*, (Marcel Dekker, 1990).

Nonpolymeric surface stabilizers are any nonpolymeric compound, such as benzalkonium chloride, a carbonium compound, a phosphonium compound, an oxonium compound, a halonium compound, a cationic organometallic compound, a quaternary phosphorous compound, a pyridinium compound, an anilinium compound, an ammonium compound, a hydroxylammonium compound, a primary ammonium compound, a secondary ammonium compound, a tertiary ammonium compound, and quaternary ammonium compounds of the formula  $NR_1R_2R_3R_4^{(+)}$ . For compounds of the formula  $NR_1R_2R_3R_4^{(+)}$ :

- (i) none of  $R_1$ - $R_4$  are  $CH_3$ ;
- (ii) one of  $R_1$ - $R_4$  is  $CH_3$ ;

- (iii) three of  $R_1$ - $R_4$  are  $CH_3$ ;
- (iv) all of  $R_1$ - $R_4$  are  $CH_3$ ;
- (v) two of  $R_1$ - $R_4$  are  $CH_3$ , one of  $R_1$ - $R_4$  is  $C_6H_5CH_2$ , and one of  $R_1$ - $R_4$  is an alkyl chain of seven carbon atoms or less;
- 5 (vi) two of  $R_1$ - $R_4$  are  $CH_3$ , one of  $R_1$ - $R_4$  is  $C_6H_5CH_2$ , and one of  $R_1$ - $R_4$  is an alkyl chain of nineteen carbon atoms or more;
- (vii) two of  $R_1$ - $R_4$  are  $CH_3$  and one of  $R_1$ - $R_4$  is the group  $C_6H_5(CH_2)_n$ , where  $n > 1$ ;
- (viii) two of  $R_1$ - $R_4$  are  $CH_3$ , one of  $R_1$ - $R_4$  is  $C_6H_5CH_2$ , and one of  $R_1$ - $R_4$  comprises at least one heteroatom;
- 10 (ix) two of  $R_1$ - $R_4$  are  $CH_3$ , one of  $R_1$ - $R_4$  is  $C_6H_5CH_2$ , and one of  $R_1$ - $R_4$  comprises at least one halogen;
- (x) two of  $R_1$ - $R_4$  are  $CH_3$ , one of  $R_1$ - $R_4$  is  $C_6H_5CH_2$ , and one of  $R_1$ - $R_4$  comprises at least one cyclic fragment;
- 15 (xi) two of  $R_1$ - $R_4$  are  $CH_3$  and one of  $R_1$ - $R_4$  is a phenyl ring; or
- (xii) two of  $R_1$ - $R_4$  are  $CH_3$  and two of  $R_1$ - $R_4$  are purely aliphatic fragments.

Such compounds include, but are not limited to, behenalkonium chloride, benzethonium chloride, cetylpyridinium chloride, behentrimonium chloride, lauralkonium chloride, cetalkonium chloride, cetrimonium bromide, cetrimonium chloride, cethylamine hydrofluoride, chlorallylmethenamine chloride (Quaternium-15), distearyldimonium chloride (Quaternium-5), dodecyl dimethyl ethylbenzyl ammonium chloride (Quaternium-14), Quaternium-22, Quaternium-26, Quaternium-18 hectorite, dimethylaminoethylchloride hydrochloride, cysteine hydrochloride, diethanolammonium POE (10) oleyl ether phosphate, diethanolammonium POE (3) oleyl ether phosphate,

25 tallow alkonium chloride, dimethyl dioctadecylammoniumbentonite, stearalkonium chloride, domiphen bromide, denatonium benzoate, myristalkonium chloride, laurtrimonium chloride, ethylenediamine dihydrochloride, guanidine hydrochloride, pyridoxine HCl, iofetamine hydrochloride, meglumine hydrochloride, methylbenzethonium chloride, myrtrimonium bromide, oleyltrimonium chloride,

30 polyquaternium-1, procaine hydrochloride, cocobetaine, stearalkonium bentonite,

stearalkoniumhectonite, stearyl trihydroxyethyl propylenediamine dihydrofluoride, tallowtrimonium chloride, and hexadecyltrimethyl ammonium bromide.

The surface stabilizers are commercially available and/or can be prepared by techniques known in the art. Most of these surface stabilizers are known pharmaceutical excipients and are described in detail in the *Handbook of Pharmaceutical Excipients*, published jointly by the American Pharmaceutical Association and The Pharmaceutical Society of Great Britain (The Pharmaceutical Press, 2000), specifically incorporated by reference.

#### 4. Nanoparticulate Angiogenesis Inhibitor/ Surface Stabilizer Particle Size

As used herein, particle size is determined on the basis of the weight average particle size as measured by conventional particle size measuring techniques well known to those skilled in the art. Such techniques include, for example, sedimentation field flow fractionation, photon correlation spectroscopy, light scattering, and disk centrifugation.

The nanoparticulate angiogenesis inhibitor compositions of the invention have an effective average particle size of less than about 2 microns. In preferred embodiments, the effective average particle size of the angiogenesis inhibitor particles is less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, or less than about 50 nm, when measured by the above techniques.

By "an effective average particle size of less than about 2000 nm" it is meant that at least 50% of the angiogenesis inhibitor particles have a particle size of less than about 2000 nm, by weight, when measured by the above techniques. Preferably, at least about 70%, about 90%, about 95%, or about 99% of the particles have a particle size of less than the effective average, *i.e.*, less than about 2000 nm, less than about 1900 nm, less than about 1800 nm, *etc.*

If the nanoparticulate angiogenesis inhibitor composition additionally comprises one or more non-angiogenesis inhibitor nanoparticulate active agents, then such active agents have an effective average particle size of less than about 2000 nm (*i.e.*, 2 microns),  
5 less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about  
10 250 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 75 nm, or less than about 50 nm, as measured by light-scattering methods, microscopy, or other appropriate methods.

If the nanoparticulate angiogenesis inhibitor is combined with a conventional or microparticulate angiogenesis inhibitor or non-angiogenesis inhibitor composition, then  
15 such a conventional composition is either solubilized or has an effective average particle size of greater than about 2 microns. By "an effective average particle size of greater than about 2 microns" it is meant that at least 50% of the conventional angiogenesis inhibitor or active agent particles have a particle size of greater than about 2 microns, by weight, when measured by the above-noted techniques. In other embodiments of the invention, at  
20 least about 70%, about 90%, about 95%, or about 99% of the conventional angiogenesis inhibitor or active agent particles have a particle size greater than about 2 microns.

## 5. Other Pharmaceutical Excipients

Pharmaceutical compositions according to the invention may also comprise one or  
25 more binding agents, filling agents, lubricating agents, suspending agents, sweeteners, flavoring agents, preservatives, buffers, wetting agents, disintegrants, effervescent agents, and other excipients. Such excipients are known in the art.

Examples of filling agents are lactose monohydrate, lactose anhydrous, and various starches; examples of binding agents are various celluloses and cross-linked  
30 polyvinylpyrrolidone, microcrystalline cellulose, such as Avicel® PH101 and Avicel®

PH102, microcrystalline cellulose, and silicified microcrystalline cellulose (ProSolv SMCC™).

Suitable lubricants, including agents that act on the flowability of the powder to be compressed, are colloidal silicon dioxide, such as Aerosil® 200, talc, stearic acid,  
5 magnesium stearate, calcium stearate, and silica gel.

Examples of sweeteners are any natural or artificial sweetener, such as sucrose, xylitol, sodium saccharin, cyclamate, aspartame, and acesulfame. Examples of flavoring agents are Magnasweet® (trademark of MAFCO), bubble gum flavor, and fruit flavors, and the like.

10 Examples of preservatives are potassium sorbate, methylparaben, propylparaben, benzoic acid and its salts, other esters of parahydroxybenzoic acid such as butylparaben, alcohols such as ethyl or benzyl alcohol, phenolic compounds such as phenol, or quarternary compounds such as benzalkonium chloride.

Suitable diluents include pharmaceutically acceptable inert fillers, such as  
15 microcrystalline cellulose, lactose, dibasic calcium phosphate, saccharides, and/or mixtures of any of the foregoing. Examples of diluents include microcrystalline cellulose, such as Avicel® PH101 and Avicel® PH102; lactose such as lactose monohydrate, lactose anhydrous, and Pharmatose® DCL21; dibasic calcium phosphate such as Emcompress®; mannitol; starch; sorbitol; sucrose; and glucose.

20 Suitable disintegrants include lightly crosslinked polyvinyl pyrrolidone, corn starch, potato starch, maize starch, and modified starches, croscarmellose sodium, cross-povidone, sodium starch glycolate, and mixtures thereof.

Examples of effervescent agents are effervescent couples such as an organic acid and a carbonate or bicarbonate. Suitable organic acids include, for example, citric,  
25 tartaric, malic, fumaric, adipic, succinic, and alginic acids and anhydrides and acid salts. Suitable carbonates and bicarbonates include, for example, sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate, magnesium carbonate, sodium glycine carbonate, L-lysine carbonate, and arginine carbonate. Alternatively, only the sodium bicarbonate component of the effervescent couple may be present.

## **6. Concentration of Nanoparticulate Angiogenesis inhibitor and Stabilizer**

The relative amount of angiogenesis inhibitor and one or more surface stabilizers can vary widely. The optimal amount of the surface stabilizers can depend, for example, upon the particular angiogenesis inhibitor selected, the hydrophilic lipophilic balance (HLB), melting point, water solubility of the surface stabilizer, and the surface tension of water solutions of the stabilizer, *etc.*

The concentration of the at least one angiogenesis inhibitor can vary from about 99.5% to about 0.001%, from about 95% to about 0.1%, or from about 90% to about 0.5%, by weight, based on the total combined weight of the at least one angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

The concentration of the one or more surface stabilizers can vary from about 0.5% to about 99.999%, from about 5.0% to about 99.9%, or from about 10% to about 99.5%, by weight, based on the total combined dry weight of the at least one angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

## **E. Methods of Making Nanoparticulate Formulations**

The nanoparticulate angiogenesis inhibitor compositions can be made using, for example, milling, precipitation, or homogenization techniques. Exemplary methods of making nanoparticulate compositions are described in the '684 patent. Methods of making nanoparticulate compositions are also described in U.S. Patent No. 5,518,187, for "Method of Grinding Pharmaceutical Substances;" U.S. Patent No. 5,718,388, for "Continuous Method of Grinding Pharmaceutical Substances;" U.S. Patent No. 5,862,999, for "Method of Grinding Pharmaceutical Substances;" U.S. Patent No. 5,665,331, for "Co-Microprecipitation of Nanoparticulate Pharmaceutical Agents with Crystal Growth Modifiers;" U.S. Patent No. 5,662,883, for "Co-Microprecipitation of Nanoparticulate Pharmaceutical Agents with Crystal Growth Modifiers;" U.S. Patent No. 5,560,932, for "Microprecipitation of Nanoparticulate Pharmaceutical Agents;" U.S. Patent No. 5,543,133, for "Process of Preparing X-Ray Contrast Compositions Containing Nanoparticles;" U.S. Patent No. 5,534,270, for "Method of Preparing Stable Drug

Nanoparticles;" U.S. Patent No. 5,510,118, for "Process of Preparing Therapeutic Compositions Containing Nanoparticles;" and U.S. Patent No. 5,470,583, for "Method of Preparing Nanoparticle Compositions Containing Charged Phospholipids to Reduce Aggregation," all of which are specifically incorporated by reference.

5 One or more non-angiogenesis inhibitor active agents can be reduced in size at the same time as the angiogenesis inhibitor, to produce a nanoparticulate angiogenesis inhibitor and nanoparticulate non-angiogenesis inhibitor active agent composition. A non-angiogenesis inhibitor active agent, which is either conventional or nanoparticulate sized, can also be added to the nanoparticulate angiogenesis inhibitor composition after  
10 particle size reduction.

In yet another embodiment of the invention, nanoparticulate angiogenesis inhibitor compositions of the invention can be made in which the formulation comprises multiple nanoparticulate angiogenesis inhibitor compositions, each of which has a different effective average particle size. Such a composition can be made by preparing the  
15 individual nanoparticulate angiogenesis inhibitor compositions using, for example, milling, precipitation, or homogenization techniques, followed by combining the different compositions to prepare a single dosage form.

The nanoparticulate angiogenesis inhibitor compositions can be utilized in solid or liquid dosage formulations, such as liquid dispersions, gels, aerosols, ointments, creams,  
20 controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, mixed immediate release and controlled release formulations, *etc.*

### **1. Milling to Obtain Nanoparticulate Dispersions**

25 Milling of aqueous angiogenesis inhibitors to obtain a nanoparticulate dispersion comprises dispersing angiogenesis inhibitor particles in a liquid dispersion medium in which the angiogenesis inhibitor is poorly soluble, followed by applying mechanical means in the presence of grinding media to reduce the particle size of the angiogenesis inhibitor to the desired effective average particle size. The angiogenesis inhibitor  
30 particles can be reduced in size in the presence of at least one surface stabilizer.

Alternatively, the angiogenesis inhibitor particles can be contacted with one or more surface stabilizers either before or after attrition. Other compounds, such as a diluent, can be added to the angiogenesis inhibitor/surface stabilizer composition either before, during, or after the size reduction process. Dispersions can be manufactured continuously or in a batch mode.

## **2. Precipitation to Obtain Nanoparticulate Angiogenesis Inhibitor Compositions**

Another method of forming the desired nanoparticulate angiogenesis inhibitor composition is by microprecipitation. This is a method of preparing stable dispersions of angiogenesis inhibitors in the presence of one or more surface stabilizers and one or more colloid stability enhancing surface active agents free of any trace toxic solvents or solubilized heavy metal impurities. Such a method comprises, for example: (1) dissolving at least one angiogenesis inhibitor in a suitable solvent; (2) adding the formulation from step (1) to a solution comprising at least one surface stabilizer to form a clear solution; and (3) precipitating the formulation from step (2) using an appropriate non-solvent. The method can be followed by removal of any formed salt, if present, by dialysis or diafiltration and concentration of the dispersion by conventional means. Dispersions can be manufactured continuously or in a batch mode.

## **3. Homogenization to Obtain Nanoparticulate Angiogenesis Inhibitor Compositions**

Exemplary homogenization methods of preparing nanoparticulate compositions are described in U.S. Patent No. 5,510,118, for "Process of Preparing Therapeutic Compositions Containing Nanoparticles." Such a method comprises dispersing angiogenesis inhibitor particles in a liquid dispersion medium, followed by subjecting the dispersion to homogenization to reduce the particle size of the angiogenesis inhibitor to the desired effective average particle size. The angiogenesis inhibitor particles can be reduced in size in the presence of at least one surface stabilizer. Alternatively, the angiogenesis inhibitor particles can be contacted with one or more surface stabilizers either before or after attrition. Other compounds, such as a diluent, can be added to the



angiogenesis inhibitor/surface stabilizer composition either before, during, or after the size reduction process. Dispersions can be manufactured continuously or in a batch mode.

5    **F.    Methods of Using Nanoparticulate Angiogenesis Inhibitor Formulations Comprising One or More Surface Stabilizers**

10    The angiogenesis inhibitor compositions of the invention are useful in treating or preventing, for example, tumor growth, cancer growth, or any mammalian disease characterized by undesirable angiogenesis.

15    The nanoparticulate compositions of the present invention can be administered to humans and animals in any pharmaceutically acceptable manner, including, but not limited to orally, pulmonary, rectally, ocularly, colonically, parenterally (*e.g.*, intravenous, intramuscular, or subcutaneous), intracisternally, intravaginally, intraperitoneally, locally (*e.g.*, powders, ointments, or drops), buccally, nasal, and topically. As used herein, the term "subject" is used to mean an animal, preferably a mammal, including a human or non-human. The terms patient and subject may be used interchangeably.

20    Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles including water, ethanol, polyols (propylene glycol, polyethylene-glycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of  
25    dispersions, and by the use of surfactants.

30    The nanoparticulate angiogenesis inhibitor compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the growth of microorganisms can be ensured by various antibacterial and antifungal agents, such as parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like.

Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, such as aluminum monostearate and gelatin.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the nanoparticulate angiogenesis inhibitor is admixed with at least one of the following: (a) one or more inert excipients (or carrier), such as sodium citrate or dicalcium phosphate; (b) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and silicic acid; (c) binders, such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; (d) humectants, such as glycerol; (e) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate; (f) solution retarders, such as paraffin; (g) absorption accelerators, such as quaternary ammonium compounds; (h) wetting agents, such as cetyl alcohol and glycerol monostearate; (i) adsorbents, such as kaolin and bentonite; and (j) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. For capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the angiogenesis inhibitor, the liquid dosage forms may comprise inert diluents commonly used in the art, such as water or other solvents, solubilizing agents, and emulsifiers. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

One of ordinary skill will appreciate that effective amounts of an angiogenesis inhibitor can be determined empirically and can be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt, ester, or prodrug form. Actual dosage levels of angiogenesis inhibitor in the nanoparticulate compositions of the invention may be varied to obtain an amount of active ingredient that is effective to obtain a desired therapeutic response for a particular composition and method of administration. The selected dosage level therefore depends upon the desired therapeutic effect, the route of

administration, the potency of the angiogenesis inhibitor, the desired duration of treatment, and other factors.

The daily dose may be administered in single or multiple doses. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the body weight, general health, sex, diet, time and route of administration, potency of the administered angiogenesis inhibitor, rates of absorption and excretion, combination with other drugs and the severity of the particular disease being treated, and like factors well known in the medical arts..

\* \* \* \* \*

The following example is given to illustrate the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in this example. Throughout the specification, any and all references to a publicly available document, including a U.S. patent, are specifically incorporated by reference.

#### **Example 1**

The purpose of this example was to describe how a nanoparticulate dispersion of an angiogenesis inhibitor can be made.

Nanocrystalline dispersions of an angiogenesis inhibitor can be made by milling the compound, at least one surface stabilizer, and any desired excipients on a suitable mill, such as a Netzsch Mill (Netzsch Inc., Exton, PA) or a Dyno-Mill, for a suitable time at a suitable temperature. 500 micron PolyMill media can be used.

#### **Example 2**

The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol, which is an angiogenesis inhibitor.

A nanoparticulate dispersion of 2-methoxyestradiol, having 5% (w/w) 2-methoxyestradiol, 1% (w/w) hydroxypropyl cellulose, low viscosity (HPC-SL), and 0.05% (w/w) docusate sodium (DOSS), was milled for 1 hour under high energy milling

conditions in a NanoMill®-001 System (Custom Machine and Design Inc., Oxford, PA; *see* U.S. Patent No. 6,431,478 for "Small Scale Mill") equipped with a 10 cc chamber and utilizing 500  $\mu$ m polymeric attrition media.

Following milling, the final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol was 153 nm, with 50% < 144 nm, 90% < 217 nm, and 95% < 251 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following two weeks storage at 5°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 195 nm.

This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor. The angiogenesis inhibitor composition having a very small effective average particle size can be sterile filtered, which is particularly useful for injectable products, and for administration to immunocompromised patients, the elderly, and infants or juveniles.

### **Example 3**

The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 5% (w/w) 2-methoxyestradiol, 1% (w/w) hydroxypropyl methylcellulose (HPMC), and 0.05% (w/w) DOSS, was milled for 1 hour under high energy milling conditions in a NanoMill®-001 System (Custom Machine and Design Inc., Oxford, PA) equipped with a 10 cc chamber and utilizing 500  $\mu$ m polymeric attrition media.

Following milling, the final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol was 162 nm, with 50% < 151 nm, 90% < 234 nm, and 95% < 277 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following two weeks storage at 5°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 193 nm.

This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### Example 4

5       The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 5% (w/w) 2-methoxyestradiol, 1% (w/w) HPC-SL, and 0.05% (w/w) DOSS, was milled for 1.5 hours under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen AG, Maschinenfabrik, Basel, Switzerland) equipped with a 150 cc batch chamber and utilizing 500  $\mu$ m polymeric attrition media.

10       The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 157 nm, with 50% < 152 nm, 90% < 212 nm, and 95% < 236 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for one month at 5°C, 25°C, and 40°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 207 nm, 216 nm, and 260 nm, respectively.

15       This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### Example 5

20       The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 5% (w/w) 2-methoxyestradiol, 1% (w/w) HPMC, and 0.05% (w/w) DOSS, was milled for 2 hours under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen AG, Maschinenfabrik, Basel, Switzerland) equipped with a 150 cc batch chamber and utilizing 500  $\mu$ m polymeric attrition media.

25       The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 157 nm, with 50% < 151 nm, 90% < 213 nm,

and 95% < 240 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for one month at 5°C, 25°C, and 40°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 182 nm, 198 nm, and 218 nm, respectively.

5           This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### **Example 6**

10           The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

          A nanoparticulate dispersion of 2-methoxyestradiol, having 15% (w/w) 2-methoxyestradiol and 4% (w/w) lysozyme was milled for 1.5 hours under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen AG, Maschinenfabrik, Basel, Switzerland) equipped with a 150 cc batch chamber and utilizing 500 µm  
15           polymeric attrition media.

          The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 110 nm, with 50% < 101 nm, 90% < 169 nm, and 95% < 204 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for one month  
20           at 5°C, 25°C, and 40°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 190 nm, 201 nm, and 202 nm, respectively.

          This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### **Example 7**

25           The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

          A nanoparticulate dispersion of 2-methoxyestradiol, having 15% (w/w) 2-methoxyestradiol, 3% (w/w) copovidonum, and 0.15% (w/w) DOSS, was milled for 1.5  
30           hours under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen

AG, Maschinenfabrik, Basel, Switzerland) equipped with a 150 cc batch chamber and utilizing 500  $\mu\text{m}$  polymeric attrition media.

The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 155 nm, with 50% < 148 nm, 90% < 217 nm, and 95% < 245 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for 1.5 months at 5°C and 25°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 209 nm and 216 nm, respectively.

This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### **Example 8**

The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 25% (w/w) 2-methoxyestradiol, 5% (w/w) HPMC, and 0.25% (w/w) DOSS, was milled for 12.6 hours under high energy milling conditions in a NanoMill®-02 System utilizing 500  $\mu\text{m}$  polymeric attrition media.

The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 149 nm, with 50% < 145 nm, 90% < 203 nm, and 95% < 223 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for one month at 5°C, 25°C, and 40°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 163 nm, 164 nm, and 167 nm, respectively.

This example demonstrates the successful preparation of a stable nanoparticulate composition of an angiogenesis inhibitor.

#### **Example 9**

The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 25% (w/w) 2-methoxyestradiol, 5% (w/w) HPMC, and 0.05% (w/w) DOSS, was milled for 3.5 hours under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen AG, Maschinenfabrik, Basel, Switzerland) equipped with a 600 cc recirculation chamber and  
5 utilizing 500  $\mu$ m polymeric attrition media.

The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 146 nm, with 50% < 143 nm, 90% < 194 nm, and 95% < 215 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). The sample showed aggregation  
10 after 4 days at 5°C and had a mean particle size of 1968 nm.

This example demonstrates that not all combinations of angiogenesis inhibitors and surface stabilizers, at all concentrations, will result in a stable nanoparticulate composition of an angiogenesis inhibitor.

#### 15 **Example 10**

The purpose of this example was to prepare a nanoparticulate composition of 2-methoxyestradiol.

A nanoparticulate dispersion of 2-methoxyestradiol, having 25% (w/w) 2-methoxyestradiol, 5% (w/w) HPMC, and 0.25% (w/w) DOSS, was milled for 5.5 hours  
20 under high energy milling conditions in a DYNO®-Mill KDL (Willy A. Bachofen AG, Maschinenfabrik, Basel, Switzerland) equipped with a 600 cc recirculation chamber and utilizing 500  $\mu$ m polymeric attrition media.

The final mean particle size (volume statistics) of the nanoparticulate dispersion of 2-methoxyestradiol following milling was 148 nm, with 50% < 144 nm, 90% < 201 nm,  
25 and 95% < 221 nm, measured using a Horiba LA-910 Laser Scattering Particle Size Distribution Analyzer (Horiba Instruments, Irvine, CA). Following storage for 4 months at 5°C, 25°C, and 40°C, the nanoparticulate dispersion of 2-methoxyestradiol had a mean particle size of 186 nm, 229 nm, and 220 nm, respectively.

This example demonstrates the successful preparation of a stable nanoparticulate  
30 composition of an angiogenesis inhibitor.



\* \* \* \*

It will be apparent to those skilled in the art that various modifications and variations can be made in the methods and compositions of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.

We claim:

1. A nanoparticulate angiogenesis inhibitor composition comprising:

- 5 (a) particles of an angiogenesis inhibitor or a salt thereof having an effective average particle size of less than about 2000 nm; and
- (b) associated with the surface thereof at least one surface stabilizer.

10 2. The composition of claim 1, wherein the angiogenesis inhibitor is selected from the group consisting of 2-methoxyestradiol, prinomastat, batimastat, BAY 12-9566, carboxyamidotriazole, CC-1088, dextromethorphan acetic, dimethylxanthenone acetic acid, EMD 121974, endostatin, IM-862, marimastat, matrix metalloproteinase,

15 penicillamine, PTK787/ZK 222584, RPI.4610, squalamine, squalamine lactate, SU5416, (+)-thalidomide, S- thalidomide, R- thalidomide, TNP-470, combretastatin, paclitaxel, tamoxifen, COL-3, neovastat, BMS-275291, SU6668, interferon-alpha, anti-VEGF antibody, Medi-522 (Vitaxin II), CAI, celecoxib, Interleukin-12, IM862, Amilloride, Angiostatin® Protein, Angiostatin K1-3, Angiostatin K1-5, Captopril, DL-alpha-

20 Difluoromethylornithine, DL-alpha-Difluoromethylornithine HCl, His-Tag® Endostatin™ Protein, Fumagillin, Herbimycin A, 4-Hydroxyphenylretinamide, gamma-interferon, Juglone, Laminin, Laminin Hexapeptide, Laminin Pentapeptide, Lavendustin A, Medroxyprogesterone, Medroxyprogesterone Acetate, Minocycline, Minocycline HCl, Placental Ribonuclease Inhibitor, Suramin, Sodium Salt Suramin, Human Platelet

25 Thrombospondin, Tissue Inhibitor of Metalloproteinase 1, Neutrophil Granulocyte Tissue Inhibitor of Metalloproteinase 1, and Rheumatoid Synovial Fibroblast Tissue Inhibitor of Metalloproteinase 2.

3. The composition of claim 1 or claim 2, wherein the angiogenesis inhibitor

30 is 2-methoxyestradiol.

4. The composition of any one of claims 1-3, wherein the angiogenesis inhibitor is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, and mixtures thereof.

5

5. The composition of any one of claims 1-5, wherein the effective average particle size of the nanoparticulate angiogenesis inhibitor is selected from the group consisting of less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.

15

6. The composition of any one of claims 1-6, wherein the composition is formulated for administration selected from the group consisting of oral, pulmonary, rectal, ophthalmic, colonic, parenteral, intracisternal, intravaginal, intraperitoneal, local, buccal, nasal, and topical administration.

20

7. The composition of any one of claims 1-6, wherein the composition is formulated into a dosage form selected from the group consisting of liquid dispersions, gels, aerosols, ointments, creams, controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, and mixed immediate release and controlled release formulations.

25

8. The composition of any one of claims 1-7, wherein the composition further comprises one or more pharmaceutically acceptable excipients, carriers, or a combination

thereof.

9. The composition of any one of claims 1-8, wherein the angiogenesis inhibitor is present in an amount selected from the group consisting of from about 99.5%  
5 to about 0.001%, from about 95% to about 0.1%, and from about 90% to about 0.5%, by weight, based on the total combined weight of the angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

10. The composition of any one of claims 1-10, wherein the at least one  
10 surface stabilizer is present in an amount selected from the group consisting of from about 0.5% to about 99.999%, from about 5.0% to about 99.9%, and from about 10% to about 99.5%, by weight, based on the total combined weight of the at least one angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

11. The composition of any one of claims 1-10, comprising at least two surface  
15 stabilizers.

12. The composition of any one of claims 1-11, wherein the surface stabilizer  
is selected from the group consisting of an anionic surface stabilizer, a cationic surface  
20 stabilizer, an ionic surface stabilizer, and a zwitterionic surface stabilizer.

13. The composition of any one of claims 1-12, wherein at least one surface  
stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin,  
casein, phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid,  
25 benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol,  
cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers,  
polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters,  
polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates,  
colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose  
30 calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose,

carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethyl-cellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; 5 poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate,  $C_{18}H_{37}CH_2C(O)N(CH_3)-CH_2(CHOH)_4(CH_2OH)_2$ , p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl  $\beta$ -D-glucopyranoside; n-decyl  $\beta$ -D-maltopyranoside; n-dodecyl 10  $\beta$ -D-glucopyranoside; n-dodecyl  $\beta$ -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl- $\beta$ -D-glucopyranoside; n-heptyl  $\beta$ -D-thioglucoside; n-hexyl  $\beta$ -D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl  $\beta$ -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- $\beta$ -D-glucopyranoside; octyl  $\beta$ -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin 15 A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

14. The composition of any one of claims 1-12, wherein the surface stabilizer is selected from the group consisting of cationic lipids, polymethylmethacrylate trimethylammonium bromide, sulfonium compounds, polyvinylpyrrolidone-2- 20 dimethylaminoethyl methacrylate dimethyl sulfate, hexadecyltrimethyl ammonium bromide, phosphonium compounds, quaternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride, 25 decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride bromide,  $C_{12-15}$ dimethyl hydroxyethyl ammonium chloride,  $C_{12-15}$ dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl 30 dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride,

lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium bromide, N-alkyl (C<sub>12-18</sub>)dimethylbenzyl ammonium chloride, N-alkyl (C<sub>14-18</sub>)dimethyl-benzyl ammonium chloride, N-tetradecylidmethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and (C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride,

5 trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyl dimethyl ammonium chloride, N-tetradecyl dimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride,

10 dodecyl dimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C<sub>12</sub> trimethyl ammonium bromides, C<sub>15</sub> trimethyl ammonium bromides, C<sub>17</sub> trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl

15 ammonium chlorides, alkyldimethylammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, POLYQUAT 10™, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearylalkonium chloride compounds, cetyl pyridinium bromide,

20 cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, MIRAPOL™, ALKAQUAT™, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

25 15. The composition of claim 14, wherein the composition is bioadhesive.

16. The composition of any one of claims 1-15, wherein the composition comprises more than one angiogenesis inhibitor.

17. The composition of claim 16, wherein at least one angiogenesis inhibitor has an effective average particle size which is greater than about 2 microns.

18. The composition of any one of claims 1-17, additionally comprising at least one nanoparticulate angiogenesis inhibitor composition having an effective average particle size of less than about 2 microns, wherein said additional nanoparticulate angiogenesis inhibitor composition has an effective average particle size which is different than the particle size of the nanoparticulate angiogenesis inhibitor composition of claim 1.

19. The composition of any one of claims 1-19, additionally comprising at least one non-angiogenesis inhibitor active agent.

20. The composition of claim 19, wherein said active agent is selected from the group consisting of amino acids, proteins, peptides, nucleotides, anti-obesity drugs, nutraceuticals, dietary supplements, central nervous symptom stimulants, carotenoids, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine, oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, alpha-adrenergic receptor blocking agents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio-pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, vasodilators, vasomodulator, xanthines, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents,

cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

21. The composition of claim 20, wherein said nutraceutical is selected from  
5 the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and  
10 probiotics.

22. The composition of any of claims 19, 20, or 21, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of less than about  
15 2 microns.

23. The composition of any of any of claims 19, 20, or 21, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of greater  
than about 2 microns.

24. The composition of any one of claims 1-23, wherein upon administration  
20 the composition redisperses such that the angiogenesis inhibitor particles have a particle size selected from the group consisting of less than about 2 microns, less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200  
25 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.



25. The composition of any one of claims 1-24, wherein the composition does not produce significantly different absorption levels when administered under fed as compared to fasting conditions.

5 26. The composition of any one of claims 1-25, wherein the difference in absorption of the nanoparticulate angiogenesis inhibitor composition of the invention, when administered in the fed versus the fasted state, is selected from the group consisting of less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 35%, less  
10 than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, and less than about 3%.

27. The composition of any one of claims 1-26, wherein the composition does not produce significantly different rates of absorption ( $T_{max}$ ) when administered under fed  
15 as compared to fasting conditions.

28. The composition of any one of claims 1-27, wherein the difference in the  $T_{max}$  for the nanoparticulate angiogenesis inhibitor composition of the invention, when administered in the fed versus the fasted state, is less than about 100%, less than about  
20 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, and less than about 3%.

29. The composition of any one of claims 1-28, wherein upon administration  
25 the  $T_{max}$  is less than that of a conventional non-nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage.

30. The composition of any one of claims 1-29, wherein in comparative pharmacokinetic testing with a non-nanoparticulate composition of the same angiogenesis  
30 inhibitor, administered at the same dosage, the nanoparticulate composition exhibits a

$T_{\max}$  selected from the group consisting of less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, and less than about 10% of the  $T_{\max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

31. The composition of any one of claims 1-30, wherein following administration the composition has a  $T_{\max}$  selected from the group consisting of less than about 2.5 hours, less than about 2.25 hours, less than about 2 hours, less than about 1.75 hours, less than about 1.5 hours, less than about 1.25 hours, less than about 1.0 hours, less than about 50 minutes, less than about 40 minutes, less than about 30 minutes, less than about 25 minutes, less than about 20 minutes, less than about 15 minutes, and less than about 10 minutes.

32. The composition of any one of claims 1-31, wherein upon administration the  $C_{\max}$  of the composition is greater than the  $C_{\max}$  of a conventional non-nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage.

33. The composition of any one of claims 1-32, wherein in comparative pharmacokinetic testing with a non-nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage, the nanoparticulate composition exhibits a  $C_{\max}$  selected from the group consisting of greater than about 5%, greater than about 10%, greater than about 15%, greater than about 20%, greater than about 30%, greater than about 40%, greater than about 50%, greater than about 60%, greater than about 70%, greater than about 80%, greater than about 90%, greater than about 100%, greater than about 110%, greater than about 120%, greater than about 130%, greater than about 140%, and greater than about 150% than the  $C_{\max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

34. A method of making an angiogenesis inhibitor composition comprising contacting particles of at least one angiogenesis inhibitor with at least one surface stabilizer for a time and under conditions sufficient to provide an angiogenesis inhibitor composition having an effective average particle size of less than about 2 microns.

5

35. The method of claim 34, wherein said contacting comprises grinding.

36. The method of claim 35, wherein said grinding comprises wet grinding.

10

37. The method of claim 34, wherein said contacting comprises homogenizing.

38. The method of claim 34, wherein said contacting comprises:

(a) dissolving the angiogenesis inhibitor particles in a solvent;

(b) adding the resulting angiogenesis inhibitor solution to a

15

solution comprising at least one surface stabilizer; and

(c) precipitating the solubilized angiogenesis inhibitor having at least one surface stabilizer associated with the surface thereof by the addition thereto of a non-solvent.

20

39. The method of any one of claims 34-38, wherein the angiogenesis inhibitor is selected from the group consisting of 2-methoxyestradiol, prinomastat, batimastat, BAY 12-9566, carboxyamidotriazole, CC-1088, dextromethorphan acetic, dimethylxanthenone acetic acid, EMD 121974, endostatin, IM-862, marimastat, matrix metalloproteinase, penicillamine, PTK787/ZK 222584, RPI.4610, squalamine, squalamine lactate, SU5416, (+)-thalidomide, S- thalidomide, R- thalidomide, TNP-470, combretastatin, paclitaxel, tamoxifen, COL-3, neovastat, BMS-275291, SU6668, interferon-alpha, anti-VEGF antibody, Medi-522 (Vitaxin II), CAI, celecoxib, Interleukin-12, IM862, Amiloride, Angiostatin® Protein, Angiostatin K1-3, Angiostatin K1-5, Captopril, DL-alpha-Difluoromethylornithine, DL-alpha-Difluoromethylornithine HCl, His-Tag® Endostatin™ Protein, Fumagillin, Herbimycin A, 4-Hydroxyphenylretinamide,

30

gamma-interferon, Juglone, Laminin, Laminin Hexapeptide, Laminin Pentapeptide, Lavendustin A, Medroxyprogesterone, Medroxyprogesterone Acetate, Minocycline, Minocycline HCl, Placental Ribonuclease Inhibitor, Suramin, Sodium Salt Suramin, Human Platelet Thrombospondin, Tissue Inhibitor of Metalloproteinase 1, Neutrophil  
5 Granulocyte Tissue Inhibitor of Metalloproteinase 1, and Rheumatoid Synovial Fibroblast Tissue Inhibitor of Metalloproteinase 2.

40. The method of any one of claims 34-39, wherein the angiogenesis inhibitor is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-  
10 crystalline phase, a semi-amorphous phase, and mixtures thereof.

41. The method of any one of claims 34-40, wherein the effective average particle size of the nanoparticulate angiogenesis inhibitor particles is selected from the group consisting of less than about 1900 nm, less than about 1800 nm, less than about  
15 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less  
20 than about 75 nm, and less than about 50 nm.

42. The method of any one of claims 34-41, wherein the angiogenesis inhibitor is present in an amount selected from the group consisting of from about 99% to about 0.001%, from about 95% to about 0.5%, and from about 90% to about 0.5%, by weight,  
25 based on the total combined weight of the angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

43. The method of any one of claims 34-42, wherein at least one surface stabilizer is present in an amount selected from the group consisting of from about 0.5%  
30 to about 99.999%, from about 5.0% to about 99.9%, and from about 10% to about 99.5%,

by weight, based on the total combined dry weight of the angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

44. The method of any one of claims 34-43, wherein the angiogenesis inhibitor particles are contacted with at least two surface stabilizers.

45. The method of any one of claims 34-44, wherein the surface stabilizer is selected from the group consisting of an anionic surface stabilizer, a cationic surface stabilizer, an ionic surface stabilizer, and a zwitterionic surface stabilizer.

46. The method of any one of claims 34-45, wherein at least one surface stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin, casein, phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethyl-cellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate,  $C_{18}H_{37}CH_2C(O)N(CH_3)CH_2(CHOH)_4(CH_2OH)_2$ , p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl  $\beta$ -D-glucopyranoside; n-decyl  $\beta$ -D-maltopyranoside; n-dodecyl  $\beta$ -D-glucopyranoside; n-dodecyl  $\beta$ -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl  $\beta$ -D-glucopyranoside; n-heptyl  $\beta$ -D-thioglucoside; n-hexyl  $\beta$ -D-glucopyranoside;

nonanoyl-N-methylglucamide; n-nonyl  $\beta$ -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- $\beta$ -D-glucopyranoside; octyl  $\beta$ -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

5

47. The method of any one of claims 34-45, wherein the surface stabilizer is selected from the group consisting of cationic lipids, polymethylmethacrylate trimethylammonium bromide, sulfonium compounds, polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate dimethyl sulfate, hexadecyltrimethyl ammonium  
 10 bromide, phosphonium compounds, quarternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl  
 15 ammonium chloride bromide, C<sub>12-15</sub>dimethyl hydroxyethyl ammonium chloride, C<sub>12-15</sub>dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium bromide, N-alkyl (C<sub>12-18</sub>)dimethylbenzyl  
 20 ammonium chloride, N-alkyl (C<sub>14-18</sub>)dimethyl-benzyl ammonium chloride, N-tetradecylidmethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and (C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride, trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium  
 25 salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyl dimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride, dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride,  
 30 lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl

benzyl dimethyl ammonium bromide, C<sub>12</sub> trimethyl ammonium bromides, C<sub>15</sub> trimethyl ammonium bromides, C<sub>17</sub> trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl ammonium chlorides, alkyl dimethyl ammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, 5 tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, POLYQUAT 10™, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearylalkonium chloride compounds, cetyl pyridinium bromide, cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, 10 MIRAPOL™, ALKAQUAT™, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

48. The method of any one of claims 34-47, wherein after preparation of a first 15 nanoparticulate angiogenesis inhibitor composition, a second angiogenesis inhibitor composition having an effective average particle size of greater than about 2 microns is combined with the first angiogenesis inhibitor composition.

49. The method of any one of claims 34-48, wherein either prior or subsequent 20 to preparation of the nanoparticulate angiogenesis inhibitor composition, at least one non-angiogenesis inhibitor active agent is added to the angiogenesis inhibitor composition.

50. The method of claim 49, wherein said non-angiogenesis inhibitor active agent is selected from the group consisting of amino acids proteins, peptides, nucleotides, 25 anti-obesity drugs, nutraceuticals, dietary supplements, carotenoids, central nervous system stimulants, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine, oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, 30 antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic

agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, alpha-adrenergic receptor blocking agents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio-pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, vasodilators, vasomodulator, xanthines, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents, cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

51. The method of claim 50, wherein said nutraceutical is selected from the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and probiotics.

52. The method of any of claims 49, 50, or 51, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of less than about 2 microns.

53. The method of any of claims 49, 50, or 51, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of greater than about 2 microns.



54. A method of treating a subject in need with an angiogenesis inhibitor composition comprising administering to the subject an effective amount of an angiogenesis inhibitor composition comprising:

- 5 (a) particles of an angiogenesis inhibitor or a salt thereof having an effective average particle size of less than about 2000 nm; and
- (b) associated with the surface thereof at least one surface stabilizer.

10 55. The method of claim 54, wherein the angiogenesis inhibitor is selected from the group consisting of 2-methoxyestradiol, prinomastat, batimastat, BAY 12-9566, carboxyamidotriazole, CC-1088, dextromethorphan acetic, dimethylxanthenone acetic acid, EMD 121974, endostatin, IM-862, marimastat, matrix metalloproteinase, penicillamine, PTK787/ZK 222584, RPI.4610, squalamine, squalamine lactate, SU5416,

15 (+)-thalidomide, S- thalidomide, R- thalidomide, TNP-470, combretastatin, paclitaxel, tamoxifen, COL-3, neovastat, BMS-275291, SU6668, interferon-alpha, anti-VEGF antibody, Medi-522 (Vitaxin II), CAI, celecoxib, Interleukin-12, IM862, Amiloride, Angiostatin® Protein, Angiostatin K1-3, Angiostatin K1-5, Captopril, DL-alpha-Difluoromethylornithine, DL-alpha-Difluoromethylornithine HCl, His-Tag®

20 Endostatin™ Protein, Fumagillin, Herbimycin A, 4-Hydroxyphenylretinamide, gamma-interferon, Juglone, Laminin, Laminin Hexapeptide, Laminin Pentapeptide, Lavendustin A, Medroxyprogesterone, Medroxyprogesterone Acetate, Minocycline, Minocycline HCl, Placental Ribonuclease Inhibitor, Suramin, Sodium Salt Suramin, Human Platelet Thrombospondin, Tissue Inhibitor of Metalloproteinase 1, Neutrophil Granulocyte Tissue

25 Inhibitor of Metalloproteinase 1, and Rheumatoid Synovial Fibroblast Tissue Inhibitor of Metalloproteinase 2.

56. The method of claim 54 or claim 55, wherein the angiogenesis inhibitor is selected from the group consisting of a crystalline phase, an amorphous phase, a semi-crystalline phase, a semi-amorphous phase, and mixtures thereof.

30

57. The method of any one of claims 54-56, wherein the effective average particle size of the nanoparticulate angiogenesis inhibitor particles is selected from the group consisting of less than about 1900 nm, less than about 1800 nm, less than about 1700 nm, less than about 1600 nm, less than about 1500 nm, less than about 1400 nm, less than about 1300 nm, less than about 1200 nm, less than about 1100 nm, less than about 1000 nm, less than about 900 nm, less than about 800 nm, less than about 700 nm, less than about 600 nm, less than about 500 nm, less than about 400 nm, less than about 300 nm, less than about 250 nm, less than about 200 nm, less than about 100 nm, less than about 75 nm, and less than about 50 nm.

58. The method of any one of claims 54-57, wherein the composition is formulated for an administration form selected from the group consisting of oral, pulmonary, rectal, ophthalmic, colonic, parenteral, intracisternal, intravaginal, intraperitoneal, local, buccal, nasal, and topical administration.

59. The method of any one of claims 54-58, wherein the composition is a dosage form selected from the group consisting of liquid dispersions, gels, aerosols, ointments, creams, controlled release formulations, fast melt formulations, lyophilized formulations, tablets, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, and mixed immediate release and controlled release formulations.

60. The method of any one of claims 54-59, wherein the composition further comprises one or more pharmaceutically acceptable excipients, carriers, or a combination thereof.

61. The method of any one of claims 54-60, wherein the angiogenesis inhibitor is present in an amount selected from the group consisting of from about 99% to about 0.001%, from about 95% to about 0.5%, and from about 90% to about 0.5%, by weight,

based on the total combined weight of the angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

62. The method of any one of claims 54-61, wherein at least one surface stabilizer is present in an amount selected from the group consisting of from about 0.5% to about 99.999%, from about 5.0% to about 99.9%, and from about 10% to about 99.5%, by weight, based on the total combined dry weight of angiogenesis inhibitor and at least one surface stabilizer, not including other excipients.

63. The method of any one of claims 54-62, wherein the angiogenesis inhibitor composition comprises at least two surface stabilizers.

64. The method of any one of claims 54-63, wherein the surface stabilizer is selected from the group consisting of an anionic surface stabilizer, a cationic surface stabilizer, an ionic surface stabilizer, and a zwitterionic surface stabilizer.

65. The method of any one of claims 54-64, wherein the at least one surface stabilizer is selected from the group consisting of cetyl pyridinium chloride, gelatin, casein, phosphatides, dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyethylene glycols, dodecyl trimethyl ammonium bromide, polyoxyethylene stearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl celluloses, hydroxypropyl methylcellulose, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethyl-cellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol, polyvinylpyrrolidone, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde, poloxamers; poloxamines, a charged phospholipid, dioctylsulfosuccinate, dialkylesters of sodium

sulfosuccinic acid, sodium lauryl sulfate, alkyl aryl polyether sulfonates, mixtures of sucrose stearate and sucrose distearate,  $C_{18}H_{37}CH_2C(O)N(CH_3)-CH_2(CHOH)_4(CH_2OH)_2$ , p-isononylphenoxypoly-(glycidol), decanoyl-N-methylglucamide; n-decyl  $\beta$ -D-glucopyranoside; n-decyl  $\beta$ -D-maltopyranoside; n-dodecyl  $\beta$ -D-glucopyranoside; n-dodecyl  $\beta$ -D-maltoside; heptanoyl-N-methylglucamide; n-heptyl-  
 5  $\beta$ -D-glucopyranoside; n-heptyl  $\beta$ -D-thioglucoside; n-hexyl  $\beta$ -D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl  $\beta$ -D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl- $\beta$ -D-glucopyranoside; octyl  $\beta$ -D-thioglucopyranoside; lysozyme, PEG-phospholipid, PEG-cholesterol, PEG-cholesterol derivative, PEG-vitamin  
 10 A, PEG-vitamin E, and random copolymers of vinyl acetate and vinyl pyrrolidone.

66. The method of any one of claims 54-64, wherein the surface stabilizer is selected from the group consisting of benzalkonium chloride, polymethylmethacrylate trimethylammonium bromide, polyvinylpyrrolidone-2-dimethylaminoethyl methacrylate  
 15 dimethyl sulfate, hexadecyltrimethyl ammonium bromide, cationic lipids, sulfonium compounds, phosphonium compounds, quaternary ammonium compounds, benzyl-di(2-chloroethyl)ethylammonium bromide, coconut trimethyl ammonium chloride, coconut trimethyl ammonium bromide, coconut methyl dihydroxyethyl ammonium chloride, coconut methyl dihydroxyethyl ammonium bromide, decyl triethyl ammonium chloride,  
 20 decyl dimethyl hydroxyethyl ammonium chloride, decyl dimethyl hydroxyethyl ammonium chloride bromide,  $C_{12-15}$ dimethyl hydroxyethyl ammonium chloride,  $C_{12-15}$ dimethyl hydroxyethyl ammonium chloride bromide, coconut dimethyl hydroxyethyl ammonium chloride, coconut dimethyl hydroxyethyl ammonium bromide, myristyl trimethyl ammonium methyl sulphate, lauryl dimethyl benzyl ammonium chloride, lauryl  
 25 dimethyl benzyl ammonium bromide, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride, lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium bromide, N-alkyl ( $C_{12-18}$ )dimethylbenzyl ammonium chloride, N-alkyl ( $C_{14-18}$ )dimethyl-benzyl ammonium chloride, N-tetradecylidmethylbenzyl ammonium chloride monohydrate, dimethyl didecyl ammonium chloride, N-alkyl and ( $C_{12-14}$ ) dimethyl 1-naphthylmethyl ammonium chloride,  
 30 trimethylammonium halide, alkyl-trimethylammonium salts, dialkyl-dimethylammonium

salts, lauryl trimethyl ammonium chloride, ethoxylated alkyamidoalkyldialkylammonium salt, an ethoxylated trialkyl ammonium salt, dialkylbenzene dialkylammonium chloride, N-didecyldimethyl ammonium chloride, N-tetradecyldimethylbenzyl ammonium, chloride monohydrate, N-alkyl(C<sub>12-14</sub>) dimethyl 1-naphthylmethyl ammonium chloride,  
5 dodecyldimethylbenzyl ammonium chloride, dialkyl benzenealkyl ammonium chloride, lauryl trimethyl ammonium chloride, alkylbenzyl methyl ammonium chloride, alkyl benzyl dimethyl ammonium bromide, C<sub>12</sub> trimethyl ammonium bromides, C<sub>15</sub> trimethyl ammonium bromides, C<sub>17</sub> trimethyl ammonium bromides, dodecylbenzyl triethyl ammonium chloride, poly-diallyldimethylammonium chloride (DADMAC), dimethyl  
10 ammonium chlorides, alkyl dimethylammonium halogenides, tricetyl methyl ammonium chloride, decyltrimethylammonium bromide, dodecyltriethylammonium bromide, tetradecyltrimethylammonium bromide, methyl trioctylammonium chloride, POLYQUAT 10™, tetrabutylammonium bromide, benzyl trimethylammonium bromide, choline esters, benzalkonium chloride, stearalkonium chloride compounds, cetyl pyridinium bromide,  
15 cetyl pyridinium chloride, halide salts of quaternized polyoxyethylalkylamines, MIRAPOL™, ALKAQUAT™, alkyl pyridinium salts; amines, amine salts, amine oxides, imide azolinium salts, protonated quaternary acrylamides, methylated quaternary polymers, and cationic guar.

20 67. The method of any one of claims 54-66, comprising additionally administering a second angiogenesis inhibitor composition having an effective average particle size of greater than about 2 microns.

25 68. The method of any one of claims 54-67, comprising additionally administering at least one non-angiogenesis inhibitor active agent.

69. The method of claim 68, wherein said non-angiogenesis inhibitor active agent is selected from the group consisting of amino acids proteins, peptides, nucleotides, anti-obesity drugs, nutraceuticals, dietary supplements, carotenoids, central nervous  
30 system stimulants, corticosteroids, elastase inhibitors, anti-fungals, alkylxanthine,

oncology therapies, anti-emetics, analgesics, opioids, antipyretics, cardiovascular agents, anti-inflammatory agents, anthelmintics, anti-arrhythmic agents, antibiotics, anticoagulants, antidepressants, antidiabetic agents, antiepileptics, antihistamines, antihypertensive agents, antimuscarinic agents, antimycobacterial agents, antineoplastic agents, immunosuppressants, antithyroid agents, antiviral agents, anxiolytics, sedatives, astringents, alpha-adrenergic receptor blocking agents, beta-adrenoceptor blocking agents, blood products, blood substitutes, cardiac inotropic agents, contrast media, corticosteroids, cough suppressants, diagnostic agents, diagnostic imaging agents, diuretics, dopaminergics, haemostatics, immunological agents, lipid regulating agents, muscle relaxants, parasympathomimetics, parathyroid calcitonin and biphosphonates, prostaglandins, radio- pharmaceuticals, sex hormones, anti-allergic agents, stimulants, anoretics, sympathomimetics, thyroid agents, vasodilators, vasomodulator, xanthines, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic analgesics, monoamine uptake inhibitors, adenosine regulating agents, cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists, and sodium channel blockers.

70. The method of claim 69, wherein said nutraceutical is selected from the group consisting of lutein, folic acid, fatty acids, fruit extracts, vegetable extracts, vitamin supplements, mineral supplements, phosphatidylserine, lipoic acid, melatonin, glucosamine/chondroitin, Aloe Vera, Guggul, glutamine, amino acids, green tea, lycopene, whole foods, food additives, herbs, phytonutrients, antioxidants, flavonoid constituents of fruits, evening primrose oil, flax seeds, fish oils, marine animal oils, and probiotics.

71. The method of any of claims 68, 69, or 70, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of less than about 2 microns.

72. The method of any of claims 68, 69, or 70, wherein at least one non-angiogenesis inhibitor active agent has an effective average particle size of greater than about 2 microns.

5 73. The method of any one of claims 54-72, wherein the composition does not produce significantly different absorption levels when administered under fed as compared to fasting conditions.

10 74. The method of any one of claims 54-73, wherein the difference in absorption of the nanoparticulate angiogenesis inhibitor composition of the invention, when administered in the fed versus the fasted state, is selected from the group consisting of less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 35%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than  
15 about 10%, less than about 5%, and less than about 3%.

75. The method of any one of claims 54-74, wherein the composition does not produce significantly different rates of absorption ( $T_{max}$ ) when administered under fed as compared to fasting conditions.

20 76. The method of any one of claims 54-75, wherein the difference in the  $T_{max}$  for the nanoparticulate angiogenesis inhibitor composition of the invention, when administered in the fed versus the fasted state, is less than about 100%, less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about  
25 50%, less than about 40%, less than about 30%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, and less than about 3%.

77. The method of any one of claims 54-76, wherein upon administration the  $T_{max}$  is less than that of a conventional non-nanoparticulate composition of the same  
30 angiogenesis inhibitor, administered at the same dosage.

78. The method of any one of claims 54-77, wherein the nanoparticulate angiogenesis inhibitor composition exhibits a  $T_{max}$ , as compared to a non-nanoparticulate composition of the same angiogenesis inhibitor administered at the same dosage, selected  
5 from the group consisting of less than about 90%, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, and less than about 10% of the  $T_{max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

10 79. The method of any one of claims 54-78, wherein upon administration the  $T_{max}$  of the composition is selected from the group consisting of less than about 2.5 hours, less than about 2.25 hours, less than about 2 hours, less than about 1.75 hours, less than about 1.5 hours, less than about 1.25 hours, less than about 1.0 hours, less than about 50  
15 minutes, less than about 40 minutes, less than about 30 minutes, less than about 25 minutes, less than about 20 minutes, less than about 15 minutes, and less than about 10 minutes.

80. The method of any one of claims 54-79, wherein upon administration the  
20  $C_{max}$  of the composition is greater than the  $C_{max}$  of a conventional non-nanoparticulate composition of the same angiogenesis inhibitor, administered at the same dosage.

81. The method of any one of claims 54-80, wherein the nanoparticulate angiogenesis inhibitor composition exhibits a  $C_{max}$ , as compared to a non-nanoparticulate  
25 composition of the same angiogenesis inhibitor administered at the same dosage, selected from the group consisting of greater than about 5%, greater than about 10%, greater than about 15%, greater than about 20%, greater than about 30%, greater than about 40%, greater than about 50%, greater than about 60%, greater than about 70%, greater than



about 80%, greater than about 90%, greater than about 100%, greater than about 110%, greater than about 120%, greater than about 130%, greater than about 140%, and greater than about 150% than the  $C_{\max}$  exhibited by the non-nanoparticulate composition of the angiogenesis inhibitor.

5

82. The method of any one of claims 54-81, wherein the method is used to treat an condition where a selective angiogenesis inhibitor is indicated.

83. The method of any one of claims 54-82, wherein the method is used to  
10 treat a mammalian disease characterized by undesirable angiogenesis.

84. The method of any one of claims 54-83, wherein the method is used to treat or prevent tumor growth.

15 85. The method of any one of claims 54-84, wherein the method is used to treat or prevent cancer growth.

86. The method of any one of claims 54-85, wherein the subject is a human.

20

# INTERNATIONAL SEARCH REPORT

national Application No  
PCT/US 03/08546

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K9/14 A61K31/565

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 24163 A (ELAN PHARMA INTERNAT LTD ;RUDDY STEPHEN B (US); RYDE NIELS P (US)) 28 March 2002 (2002-03-28) cited in the application page 9, line 18 - line 20 example 4; tables 5,6	1,2,4-86
X	EP 0 499 299 A (STERLING WINTHROP INC) 19 August 1992 (1992-08-19) cited in the application page 3, line 19 - line 38	1,2,4-86
Y	page 3, line 54 -page 4, line 9 page 4, line 16 -page 5, line 32 page 13, line 48 - line 50 examples See p.4, 1.5: tamoxifen, medroxyprogesterone	1-86
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

9 July 2003

Date of mailing of the international search report

17/07/2003

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Giménez Miralles, J

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/08546

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 577 215 A (STERLING WINTHROP INC) 5 January 1994 (1994-01-05) cited in the application page 2, line 45 -page 3, line 26	1,2,4-86
Y	page 3, line 49 -page 4, line 9 page 4, line 37 - line 58 examples 11-16 See p.3, 1.9 ff: taxol, medroxyprogesterone, ethynylestradiol, tamoxifen, thalidomide ---	1-86
Y	US 2002/002294 A1 (CUSHMAN MARK ET AL) 3 January 2002 (2002-01-03) page 1, paragraphs 5,7 page 3, paragraphs 29,37 page 4, paragraphs 45,46; examples 4,5; tables 1,2 See tables 1 and 2: 2-methoxyestradiol, 17-ethynylestradiol ---	3
Y	WO 00 53164 A (LIVERSIDGE ELAINE ;NANOSYSTEMS (US); WEI LINDEN (US); GOTTARDY GRE) 14 September 2000 (2000-09-14) cited in the application page 11 -page 12 claim 1; examples the whole document ---	1-86
Y	US 6 068 858 A (LIVERSIDGE GARY G ET AL) 30 May 2000 (2000-05-30) cited in the application column 7, line 25 - line 32 column 7, line 66 -column 8, line 28 examples 8,9 ---	1-86
Y	WO 01 26635 A (ELAN PHARMA INTERNAT LTD ;BOSCH H WILLIAM (US); COOPER EUGENE R (U) 19 April 2001 (2001-04-19) cited in the application page 23, line 6 -page 26, line 11 examples ---	1-86
Y	US 5 510 118 A (SWANSON JON R ET AL) 23 April 1996 (1996-04-23) cited in the application column 5, line 39 - line 41 claim 1; table I ---	1-86
Y	WO 90 15593 A (YTKEMISKA INST) 27 December 1990 (1990-12-27) page 3, line 18 - line 34 page 5, line 22 page 7, line 29 examples -----	1-86

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.1

Although claims 54-86 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

## Continuation of Box I.1

Claims Nos.: 54-86

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

-----

## Continuation of Box I.2

Present independent claims 1 and 34 relate to an extremely large number of possible compositions (combination of an "angiogenesis inhibitor" and a "surface stabilizer", any combination being possible with no particular limitation). Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compositions claimed, namely those wherein the angiogenesis inhibitor is 2-methoxyestradiol, for the following reasons:

The technical problem to be solved is the provision of stable nanoparticulate dispersions of angiogenesis inhibitors comprising a surface stabilizer wherein the nanoparticles (having a mean particle size of ca. 200 nm) do not aggregate/agglomerate significantly following storage, i.e. the mean particle size of the nanoparticulate dispersions does not significantly grow upon storage. The problem of particle aggregation in nanoparticulate dispersions depends on surface interactions between the particles. It is obvious that such surface interactions will be different for different active agents and different surface stabilizers as a function of their physicochemical characteristics. Not all surface stabilizers will function to produce a non-agglomerated nanoparticulate composition for all active agents. The application provides data showing that the relevant technical problem is solved for 2-methoxyestradiol by means of stable nanocrystalline dispersions of said compound and a surface stabilizer such as HPC, HPMC, lysozyme, or copovidonum (see the examples). However, claim 2 recites a large list of other possible angiogenesis inhibitors, and claims 13 and 14 recite a large list of other possible surface stabilizers, having very different physicochemical properties. The content of the application does not provide any evidence at all in order to make credible that the relevant technical problem is also solved for each one of the possible angiogenesis inhibitor-surface stabilizer combinations claimed. Therefore, nanoparticulate dispersions of the angiogenesis inhibitors recited in claim 2 other than 2-methoxyestradiol are not considered to be

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

sufficiently disclosed in the application.

Accordingly, in the present case claims 1 and 34 so lack support, and/or the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and sufficiently disclosed, namely those parts relating to the compositions wherein the angiogenesis inhibitor is 2-methoxyestradiol in the sense of claim 3.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

international application No.  
PCT/US 03/08546

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 54-86  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: -  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 03/08546

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0224163	A	28-03-2002	US 6375986 B1	23-04-2002
			AU 9501701 A	02-04-2002
			CA 2416109 A1	28-03-2002
			EP 1318788 A1	18-06-2003
			WO 0224163 A1	28-03-2002
			US 2002110597 A1	15-08-2002
			US 2003108616 A1	12-06-2003
EP 0499299	A	19-08-1992	US 5145684 A	08-09-1992
			AT 184202 T	15-09-1999
			AT 195416 T	15-09-2000
			AU 642066 B2	07-10-1993
			AU 1014592 A	30-07-1992
			AU 654836 B2	24-11-1994
			AU 1014792 A	30-07-1992
			CA 2059431 A1	26-07-1992
			CA 2059432 A1	26-07-1992
			DE 69229925 D1	14-10-1999
			DE 69229925 T2	17-02-2000
			DE 69231345 D1	21-09-2000
			DE 69231345 T2	26-04-2001
			DK 498482 T3	27-03-2000
			DK 499299 T3	02-01-2001
			EP 0498482 A2	12-08-1992
			EP 0499299 A2	19-08-1992
			ES 2139586 T3	16-02-2000
			ES 2149164 T3	01-11-2000
			FI 920321 A	26-07-1992
			FI 920322 A	26-07-1992
			GR 3032059 T3	31-03-2000
			GR 3034759 T3	28-02-2001
			HU 62462 A2	28-05-1993
			HU 60635 A2	28-10-1992
			IE 920217 A1	29-07-1992
			IE 920218 A1	29-07-1992
			IL 100754 A	16-10-1996
			IL 100755 A	08-12-1995
			JP 4317053 A	09-11-1992
			JP 4295420 A	20-10-1992
			KR 200061 B1	15-06-1999
			MX 9200291 A1	01-10-1992
			MX 9200292 A1	01-10-1992
			NO 920333 A	27-07-1992
			NO 920334 A	27-07-1992
			NZ 241361 A	25-06-1993
			NZ 241362 A	25-06-1993
			PT 499299 T	31-01-2001
			SG 55104 A1	21-12-1998
			RU 2074002 C1	27-02-1997
			RU 2066553 C1	20-09-1996
			US 5451393 A	19-09-1995
			US 5494683 A	27-02-1996
			US 5552160 A	03-09-1996
			US 5399363 A	21-03-1995
			US 5318767 A	07-06-1994
EP 0577215	A	05-01-1994	US 5399363 A	21-03-1995
			AT 190835 T	15-04-2000

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/08546

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0577215 A		AU 675432 B2	06-02-1997
		AU 4156093 A	06-01-1994
		CA 2098242 A1	02-01-1994
		CN 1084391 A ,B	30-03-1994
		CZ 9301316 A3	16-02-1994
		DE 69328136 D1	27-04-2000
		DE 69328136 T2	09-11-2000
		DK 577215 T3	28-08-2000
		EP 0577215 A1	05-01-1994
		ES 2143488 T3	16-05-2000
		FI 933040 A	02-01-1994
		HU 64832 A2	28-03-1994
		JP 7165562 A	27-06-1995
		MX 9303950 A1	31-01-1994
		NO 932403 A	03-01-1994
		NZ 248042 A	26-10-1994
		RU 2130781 C1	27-05-1999
		SG 55089 A1	21-12-1998
		SK 68193 A3	02-02-1994
		US 5494683 A	27-02-1996
US 2002002294 A1	03-01-2002	NONE	
WO 0053164 A	14-09-2000	US 6267989 B1	31-07-2001
		AT 227983 T	15-12-2002
		AU 3230600 A	28-09-2000
		CA 2367096 A1	14-09-2000
		DE 60000823 D1	02-01-2003
		EP 1161229 A1	12-12-2001
		JP 2002538199 A	12-11-2002
		WO 0053164 A1	14-09-2000
US 6068858 A	30-05-2000	WO 9835666 A1	20-08-1998
		US 6221400 B1	24-04-2001
		US 6045829 A	04-04-2000
WO 0126635 A	19-04-2001	US 2002068092 A1	06-06-2002
		AU 7993800 A	23-04-2001
		CA 2386782 A1	19-04-2001
		EP 1217993 A2	03-07-2002
		JP 2003511406 T	25-03-2003
		WO 0126635 A2	19-04-2001
US 5510118 A	23-04-1996	AU 4867396 A	04-09-1996
		WO 9625152 A1	22-08-1996
WO 9015593 A	27-12-1990	SE 464743 B	10-06-1991
		AU 5937590 A	08-01-1991
		SE 8902257 A	22-12-1990
		WO 9015593 A1	27-12-1990